

An update on nuclear calcium signalling

Martin D. Bootman¹, Claire Fearnley¹, Ioannis Smyrniias¹, Fraser MacDonald² and H. Llewelyn Roderick^{1,3,*}

¹Laboratory of Molecular Signalling, The Babraham Institute, Babraham, Cambridge CB22 3AT, UK

²KCL Dental Institute, Department of Orthodontics, St Thomas Street, London SE1 9RT, UK

³Department of Pharmacology, University of Cambridge, Tennis Court Road, Cambridge CB2 1PD, UK

*Authors for correspondence (e-mails: martin.bootman@bbsrc.ac.uk; llewelyn.roderick@bbsrc.ac.uk)

Journal of Cell Science 122, 2337–2350 Published by The Company of Biologists 2009
doi:10.1242/jcs.028100

Summary

Over the past 15 years or so, numerous studies have sought to characterise how nuclear calcium (Ca^{2+}) signals are generated and reversed, and to understand how events that occur in the nucleoplasm influence cellular Ca^{2+} activity, and vice versa. In this Commentary, we describe mechanisms of nuclear Ca^{2+} signalling and discuss what is known about the origin and physiological significance of nuclear Ca^{2+} transients. In particular, we focus on the idea that the nucleus has an

autonomous Ca^{2+} signalling system that can generate its own Ca^{2+} transients that modulate processes such as gene transcription. We also discuss the role of nuclear pores and the nuclear envelope in controlling ion flux into the nucleoplasm.

Key words: Calcium, Gene transcription, InsP_3 , Nucleus, Paclitaxel, Ryanodine, Signalling, Spindle checkpoint, Taxol

Introduction

Calcium (Ca^{2+}) is a key player in signal transduction. It modulates diverse cellular activities ranging from fertilisation to cell death (Berridge et al., 2000). It is well known that different cell types use various elements from a broad Ca^{2+} signalling ‘toolkit’ (Box 1); consequently, the characteristics of global Ca^{2+} signals, and their physiological effects, can vary considerably (Berridge et al., 2000). This might also be the case for nuclear Ca^{2+} signalling, which is essentially a complex form of local Ca^{2+} signalling. As described below, the mechanisms of nuclear Ca^{2+} signalling are extensive, and it is likely that there is considerable flexibility in how Ca^{2+} signals are triggered in the nucleus and in the downstream targets that are affected. In this Commentary, we discuss what is known about the origin and physiological significance of nuclear Ca^{2+} transients. In particular, we focus on the idea that the nucleus has an autonomous Ca^{2+} signalling system that can generate its own Ca^{2+} transients, which modulate processes such as gene transcription. We also discuss the role of nuclear pores and the nuclear envelope in controlling ion flux into the nucleoplasm. It should be noted that nuclear Ca^{2+} signalling has been the subject of previous reviews, some of which have presented different points of view (Bootman et al., 2000; Gerasimenko and Gerasimenko, 2004; Gomes et al., 2006; Lee et al., 1998; Santella and Carafoli, 1997).


The physiological significance of nuclear Ca^{2+} signalling

It has been well established that Ca^{2+} signals occurring inside cells affect activities within the nucleus. For example, the amplitude and frequency of global cellular Ca^{2+} signals can regulate the transcription of various genes (Dolmetsch et al., 1998). But what is the significance of the nuclear component of these Ca^{2+} transients? Do patterns of nuclear Ca^{2+} signalling simply track cytosolic Ca^{2+} signalling, or are there independent nuclear Ca^{2+} signalling mechanisms that are necessary for proper cell function?

A precise method for analysing the function of nuclear Ca^{2+} is to specifically localise a buffering molecule within the nucleoplasm, which prevents changes in nuclear Ca^{2+} levels but does not affect cytosolic Ca^{2+} signals. Nathanson and colleagues employed this

approach to examine the involvement of nuclear Ca^{2+} in regulating the proliferation of hepatocyte cell lines (Rodrigues et al., 2007). They observed that targeting the Ca^{2+} -binding protein parvalbumin to the nucleus significantly reduced cell proliferation and altered the proportions of cells that were in the S and G₂-M phases of the cell cycle. By contrast, the expression of parvalbumin in the cytosol did not alter cell proliferation or cell cycle distribution. Most interestingly, the expression of the parvalbumin in the nucleus, but not in the cytosol, reduced the development of tumours when HepG2 cells were subcutaneously implanted into nude mice. Using a similar approach, the nuclear expression of parvalbumin was found to inhibit activation of the Elk1 transcription factor following stimulation of hepatoma cells with epidermal growth factor (EGF), whereas buffering cytosolic Ca^{2+} did not have the same effect (Pusl et al., 2002). Collectively, these data suggest that nuclear Ca^{2+} signals are important for regulating cell proliferation in some contexts.

Ca^{2+} is a central player in many cellular signal-transduction cascades that modulate gene transcription. Signalling cascades mediate changes in gene expression by stimulating the translocation of transcription factors from the cytosol to the nucleus, or by causing the translocation and/or activation of enzymes that regulate the activity of nuclear transcription factors or the structure of chromatin. Although an increase in the concentration of nuclear Ca^{2+} might not be the initiating signal for such events, it can be necessary for gene transcription to occur or to be sustained. For example, nuclear Ca^{2+} can activate gene transcription via the nuclear factor of activated T cells (NFAT) family of proteins. In quiescent cells, NFAT is mainly found in the cytosol, but in response to an increase in cytosolic Ca^{2+} , it rapidly (~10 minutes) translocates to the nucleus (Tomida et al., 2003). The translocation of NFAT occurs following its dephosphorylation by the Ca^{2+} -calmodulin-sensitive phosphatase calcineurin (Mellstrom et al., 2008), which exposes an intrinsic nuclear localisation sequence (NLS) in NFAT. NFAT and calcineurin associate in a Ca^{2+} -dependent manner, and it has been reported that they translocate into the nucleus as a complex (Shibasaki et al., 1996). When Ca^{2+} is elevated in the nucleus, calcineurin remains active and associated with NFAT, and thereby sustains its transcriptional activity. Calcineurin mediates this effect by outcompeting kinases that promote NFAT re-phosphorylation

Box 1. Cellular Ca^{2+} -signalling machinery

Cytosolic Ca^{2+} signals can be evoked by chemical stimuli (such as hormones, growth factors and toxins), environmental changes (such as changes in pH or temperature shifts), mechanical deformation or depolarisation. Depending on the cell type, the nature of the stimulus and the extent of stimulation, Ca^{2+} signals can be transient, oscillatory or sustained (Thomas et al., 1997), and can occur globally or as subcellular events (Bootman et al., 2001). To generate such a diverse spectrum of signals, cells employ a range of messengers and mechanisms to evoke Ca^{2+} fluxes between various cellular compartments. For example, cells can access Ca^{2+} from a variety of intracellular stores (including the sarcoplasmic or endoplasmic reticulum, the Golgi, acidic vesicles and the NE) and via influx from the extracellular space. The figure depicts the basic Ca^{2+} cycle present in all mammalian cell types. The pathways that increase cytosolic Ca^{2+} levels are shown with red arrows and the mechanisms that reduce Ca^{2+} levels are depicted with blue arrows.

Ca^{2+} influx can occur via several different types of channels that have diverse activation mechanisms. The characteristics of the Ca^{2+} signals evoked by these channels depends on their biophysical properties, their expression levels and their location within the plasma membrane of the cells. Release of Ca^{2+} from intracellular stores is also mediated by several different types of Ca^{2+} channel, of which InsP_3Rs (see Box 2) and RyRs are the best characterised. The magnitude of Ca^{2+} signals can be limited by various buffers inside cells, such as Ca^{2+} -binding proteins and mitochondria. These buffers shape the spatial domain, time-course and amplitude of Ca^{2+} rises. Elevation of cytosolic Ca^{2+} concentration is transduced into functional changes in cellular activity by numerous sensors that bind Ca^{2+} . Cellular Ca^{2+} sensors have varying affinities and locations, allowing them to specifically respond to different Ca^{2+} patterns.

(which would cause its nuclear export) and through a physical interaction that occludes a nuclear-export sequence (NES) in the NFAT protein.

An example of a nuclear-localised transcription factor that is regulated by both nuclear and cytosolic Ca^{2+} signals is cyclic AMP response element-binding protein (CREB). The signalling mechanisms that underlie CREB activation have been worked out particularly well in neurons, in which it has been established that depolarisation leads to the rapid phosphorylation of CREB on Ser133 by Ca^{2+} -calmodulin-dependent kinase IV, which provides a necessary, but not sufficient, signal for CREB-mediated gene transcription (Dolmetsch et al., 2001). The triggering event is an

increase in cytosolic Ca^{2+} levels in the immediate vicinity of L-type Ca^{2+} channels or N-methyl-D-aspartate receptors that are found in the synapse, at a site that is distal to the nucleus (Shaywitz and Greenberg, 1999). As CREB is only found in the nucleus, the phosphorylation of Ser133 is mediated by one of several Ca^{2+} -sensitive signal transduction cascades that convey information from the remote synapses into the nucleus. In the nucleus, phosphorylated CREB binds additional proteins to form a transcriptionally active complex (Shaywitz and Greenberg, 1999). Although the synaptic Ca^{2+} signal does not need to propagate to the nucleus to induce phosphorylation of Ser133 on CREB, an increase in the level of nuclear Ca^{2+} is required for CREB-mediated gene transcription to occur. This is because additional Ca^{2+} -dependent phosphorylation of CREB and its co-activators is required (Chawla et al., 1998; Kornhauser et al., 2002). Indeed, nuclear injection of an artificial Ca^{2+} buffer prevents CREB-mediated gene transcription, but not transcription induced by the serum-response element, which is sensitive to cytosolic Ca^{2+} buffering (Hardingham et al., 1997).

Another well-known target of nuclear Ca^{2+} signalling is the transcriptional regulator downstream regulatory element antagonist modulator (DREAM). It is well established that DREAM represses the expression of prodynorphin, an opiate-receptor precursor, and that mice that lack DREAM have a constitutive analgesic condition (Cheng et al., 2002). DREAM contains four Ca^{2+} -binding EF hands (a widely expressed structural motif in proteins that bind Ca^{2+}), and is directly regulated by an increase in nuclear Ca^{2+} levels (Mellstrom et al., 2008). When the concentration of Ca^{2+} in the nucleus is low, DREAM represses gene transcription by binding to DNA and displacing other transcriptional activators such as CREB (Ledo et al., 2002). An increase in the concentration of nuclear Ca^{2+} causes DREAM to dissociate from its DNA binding sites and thereby allows the transcription of its target genes.

Space limitations prevent a comprehensive discussion of the downstream targets of Ca^{2+} signalling pathways in the nucleus. However, it is evident from the *in vitro* and *in vivo* experiments described above that nucleoplasmic and cytosolic Ca^{2+} signals can synergistically control important cellular events. Furthermore, the nuclear component of global Ca^{2+} signals – or an independent increase in nuclear Ca^{2+} alone – can regulate specific nuclear processes.

Mechanisms that regulate the level or function of Ca^{2+} in the nucleus

Changes in nuclear Ca^{2+} concentration clearly impact on many cellular functions, but what are the mechanisms involved in regulating nuclear Ca^{2+} levels, and how are they related to those that operate in the cytosol? In the following section, we describe the mechanisms by which Ca^{2+} moves into and out of the nucleus and how intranuclear Ca^{2+} stores are triggered. Together, these mechanisms contribute to the regulation of Ca^{2+} -dependent nuclear signalling pathways (Fig. 1).

InsP_3Rs , RyRs and NAADPRs

To date, the most widely implicated mechanism for nuclear Ca^{2+} release is via the activation of inositol(1,4,5)-trisphosphate receptors (InsP_3Rs) (see Box 2). Evidence in support of a role for nuclear InsP_3Rs in regulating nuclear Ca^{2+} release has been obtained by using InsP_3 -binding assays (Humbert et al., 1996), InsP_3R -specific antibodies (Leite et al., 2003; Malviya, 1994; Stehno-Bittel et al., 1995), electrophysiological recordings from nuclear envelope (NE) patches or from whole nuclei (Bustamante et al., 2000; Stehno-Bittel




Fig. 1. Summary of the pathways that regulate nuclear Ca^{2+} levels. Key aspects of this schematic are the separate Ca^{2+} -releasing mechanisms in the cytosol and nucleus. In particular, the independent cytosolic and nuclear phosphoinositide cycles are depicted. Although Ca^{2+} release and the production of Ca^{2+} -releasing messengers can be localised in the individual compartments, nuclear and cytosolic Ca^{2+} levels can also be coordinated via the diffusion of messengers through NPCs. Some of the putative pathways described in this figure are controversial. In particular, the translocation of growth-factor receptors into the nucleus has not been observed in all studies (marked by '?'). + and – indicate a positive or negative effect on a downstream process, respectively.

et al., 1995), nuclear microinjection of InsP_3 (Hennager et al., 1995; Huh et al., 2007; Santella et al., 2003), direct visualisation of InsP_3 -evoked nuclear Ca^{2+} signals (Higazi et al., 2009), reconstitution of nuclear InsP_3 Rs into lipid bilayers (Leite et al., 2003) and prevention of nuclear Ca^{2+} signals by using InsP_3 R antagonists (Kumar et al., 2008). Furthermore, the addition of InsP_3 to nuclei that were isolated from *Xenopus* oocytes, *Aplysia* neurons, mammalian epithelial cells or pancreatic acinar cells (Bezin et al., 2008a; Gerasimenko et al., 1995; Quesada and Verdugo, 2005; Stehno-Bittel et al., 1995) has been shown to induce nucleoplasmic Ca^{2+} transients. Not all of these studies differentiated between expression of InsP_3 Rs on the inner or outer NE. However, it is clear that InsP_3 Rs are present on both the inner and outer NE, and that those channels expressed on the inner NE can trigger the release of Ca^{2+} from the NE lumen directly into the nucleoplasm. The actual proportion of the total cellular InsP_3 Rs that are found on the inner NE (or other putative nuclear stores; see below) is probably very small (<1%) (Laflamme et al., 2002). However, their privileged access to the nucleoplasm might confer them with an essential local signalling function.

It has also been proposed that InsP_3 Rs are present within heterochromatin and euchromatin, where they are expressed on small vesicular Ca^{2+} stores that also contain high-capacity Ca^{2+} -binding chromogranin proteins (Yoo et al., 2007). The presence of small InsP_3 -sensitive vesicular stores could help to explain the curious spotted distribution of nuclear InsP_3 R staining that is sometimes observed when using InsP_3 R-specific antibodies (particularly type-2- InsP_3 R-specific antibodies) (Garcia et al., 2004; Laflamme et al., 2002; Leite et al., 2003). If this is the case, these vesicular structures could represent nuclear InsP_3 -sensitive Ca^{2+}


stores that might operate independently of cytosolic Ca^{2+} channels. Exactly how these vesicles are formed and how chromogranins and InsP_3 Rs are trafficked to them is unclear. It is known that chromogranins can interact with InsP_3 Rs and potentiate their opening (Thrower et al., 2003), and it has been proposed that the colocalisation of chromogranins with InsP_3 Rs on nuclear vesicles underlies the increased sensitivity of nuclei to InsP_3 -evoked Ca^{2+} release (Huh et al., 2007). However, more work is needed to verify the existence and function of these nucleoplasmic Ca^{2+} stores.

In addition to InsP_3 Rs, it has been demonstrated that ryanodine receptors (RyRs) and nicotinic acid adenine dinucleotide phosphate receptors (NAADPRs) are expressed inside, or on the NE, of nuclei in various cell types, and that they regulate nuclear Ca^{2+} signalling. For example, cyclic ADP ribose or NAADP have been shown to cause the release of Ca^{2+} from the NE and to evoke Ca^{2+} transients in nuclei isolated from *Aplysia* neurons (Bezin et al., 2008b), osteoblasts (Adebanjo et al., 2000) and pancreatic acinar cells (Gerasimenko et al., 2003).

The nucleoplasmic reticulum

Ca^{2+} -release channels have been found on nucleoplasm-facing invaginations of the NE, which are known as the nucleoplasmic reticulum (or nuclear tubules) (Echevarria et al., 2003; Lee et al., 2006; Lui et al., 2003; Schermelleh et al., 2008). Although such structures have been observed in a wide range of cell types (Fricker et al., 1997), they might not be universal (Bezin et al., 2008b). For reasons that are unknown, the structure of the nucleoplasmic reticulum seems to vary from being made up of simple solitary projections (Fig. 2) to a complex branched network, and it is interesting that the tubules often colocalise with nucleoli (Fricker

Box 2. InsP_3 Rs are a generic pathway for Ca^{2+} release



With respect to Ca^{2+} release from intracellular stores, the best-characterised mechanisms involve the activation of InsP_3 Rs, RyRs and NAADPRs. InsP_3 is a highly diffusible hydrophilic messenger that is produced by the hydrolysis of $\text{PtdIns}(4,5)\text{P}_2$ (Irvine, 2006). This is catalysed by a family of PLC isozymes, which, depending on the isozyme, can be activated by G-proteins (Gq), phosphorylation or Ca^{2+} itself, or are introduced into egg cytosol during fertilisation (see figure). The other product of $\text{PtdIns}(4,5)\text{P}_2$ hydrolysis is DAG, which stays within the plasma membrane where it promotes the activation of protein kinase C or is metabolised further (some of the metabolites promote Ca^{2+} signals). InsP_3 -induced Ca^{2+} release is probably the most widespread mechanism for Ca^{2+} mobilisation in mammalian cells, and is a focal point for the convergence of multiple signalling inputs (Roderick and Bootman, 2003).

RyRs are structurally similar to InsP_3 Rs in that they comprise a tetrameric pinwheel-like arrangement, but have approximately twice the molecular mass and conductance as InsP_3 Rs. They have a more restricted expression profile than InsP_3 Rs (Bennett et al., 1996), and are best characterised in their role as Ca^{2+} -induced Ca^{2+} -release channels in muscle and neurons. The opening of RyRs can be modulated by cADPR, which is produced from NAD^+ by the action of ribosyl cyclase enzymes such as CD38 (Bezin et al., 2008a). The same enzymes may also catalyse an alternative reaction that culminates in the production of a different Ca^{2+} -releasing messenger, NAADP (Soares et al., 2007). In addition to InsP_3 , cADPR and NAADP, Ca^{2+} mobilisation can also be evoked by metabolites including nitric oxide, DAG, arachidonic acid, sphingosine and sphingosine-1-phosphate and Ca^{2+} itself (Bootman et al., 2002).

et al., 1997). The lumen in the centre of the nucleoplasmic reticulum tubules is contiguous with the cytosol and presumably allows cytosolic messengers to gain increased access to the deeper parts of the nucleus. There is evidence that functional Ca^{2+} -release channels localise on the nucleoplasmic reticulum (Echevarria et al., 2003; Marius et al., 2006). For example, photorelease of caged InsP_3 (Echevarria et al., 2003) or caged Ca^{2+} (Marius et al., 2006) in the vicinity of the nucleoplasmic reticulum initiated regenerative nuclear Ca^{2+} signals. These data indicate that InsP_3 Rs and RyRs are present on the nucleoplasmic reticulum. It has been suggested that the presence of a nucleoplasmic reticulum increases the




Fig. 2. An example of nucleoplasmic reticulum. (A) Single HeLa cell expressing ER-targeted EGFP. The net-like structure of the ER can be seen, especially at the periphery of the cell where the tubules of the organelle are less dense. The position of the nucleus is marked by N. (B) Confocal sections of the cell were volume-rendered to provide a 3D image of the fluorescence labelling. This 3D image was then serially sectioned in the vertical plane to produce the images in Bi-v. The images in Bi-v therefore represent optical sections running from the top to the bottom of the cell. An example of a position from which a vertical section was obtained is indicated by the blue bar in A. The nucleus is evident as an unlabelled region in the middle of the cell. Biii-v indicate a membranous protrusion that spans across the nucleus from one side to the other, which represents the nucleoplasmic reticulum. Images were obtained using using a Zeiss LSM 510 laser-scanning confocal microscope. The optical depth of each image is approximately 1 μm.

surface-to-volume ratio within the nucleus, thereby facilitating both the entry and exit of Ca^{2+} .

Calmodulin

As indicated above, cytosolic and nuclear activities can be coordinated by the spread of Ca^{2+} between these compartments. In

addition, the Ca^{2+} -dependent movement of enzymes (such as calcineurin) or other signalling mediators can convey signals into and out of the nucleus. One such moiety is the ubiquitous Ca^{2+} -binding protein calmodulin, which has been proposed to translocate into the nucleus following cellular stimulation. Although translocation of calmodulin from the cytosol into the nucleus has been observed in numerous cell types following elevation of cytosolic Ca^{2+} (Mermelstein et al., 2001; Thorogate and Torok, 2004; Wu and Bers, 2007), the mechanism by which calmodulin enters the nucleus is not clear. Calmodulin does not contain an obvious motif for nuclear localisation, and might therefore 'piggy-back' on other proteins that are translocated into the nucleus (Thorogate and Torok, 2007).

Although calmodulin is constitutively present in the nucleus, translocation of additional calmodulin, together with an increase in Ca^{2+} levels, could be crucial for nuclear signalling pathways. For example, calmodulin translocation in depolarised neurons occurred over the same time-course as CREB phosphorylation (Deisseroth et al., 1998). Buffering nuclear calmodulin, or inhibiting the nuclear influx of calmodulin, prevented the depolarisation-induced phosphorylation of CREB on Ser133. Furthermore, transgenic mice expressing a calmodulin-binding protein that was specifically expressed in the nuclei of forebrain neurons had a decreased level of CREB phosphorylation and impaired long-term memory consolidation (Limback-Stokin et al., 2004). Calmodulin is one example of many proteins that probably move into the nucleus to allow Ca^{2+} signals to function. Ca^{2+} causes its many functions by binding to proteins and altering their configurations to make them active or inactive. Therefore, the movement of signalling proteins concomitant with nucleoplasmic Ca^{2+} changes is important for Ca^{2+} to trigger downstream events.


Can NPCs inhibit Ca^{2+} diffusion?

Nuclear pore complexes (NPCs) are the major gateway for ions and macromolecules to pass between the cytosol and nucleoplasm (Box 3). Measurements of the size of the NPC central channel indicate that particles smaller than 9 nm should pass through without restriction. In particular, Ca^{2+} ions, which have a hydrated radius of ~ 4 Å, should pass through easily. Although the central NPC channel can become partially occluded during the transport of various cargos, which could plausibly affect Ca^{2+} movement, ions and small molecules can pass unhindered through other channels that are thought to exist at the periphery of NPCs (Kramer et al., 2007). From these simple considerations, it would be expected that NPCs provide a constitutive conduit for Ca^{2+} moving between the nucleoplasm and the cytosol.

However, it has been suggested that NPCs can attain a conformation in which they do not permit ion transport. Using patch-clamp techniques, it is possible to form giga-ohm seals on the outer NE for periods of many minutes without any evident ion flux (Bustamante et al., 2000). This is a striking finding, as even a conservative estimate of NPC density suggests that a typical patch-clamp experiment should encompass 10–100 NPCs. As the pores have a conductance of ~ 1 nanosiemen each, there should be a considerable current whenever a patch electrode is attached (Danker et al., 1997). However, electrophysiological recordings often detect only a few (about six per patch) operative channels (Bustamante et al., 2000). These observations have led to the suggestion that NPCs mainly exist in a closed conformation in which they are not ion conductive.

An alternative viewpoint has arisen from studies carried out using a different electrophysiological approach to monitor NPC activity.

Box 3. Nuclear pore complexes



NPCs are large (~ 125 MDa) protein complexes with an eightfold-symmetric architecture. They are approximately 120 nm in diameter, 80 nm in height and are studded around the circumference of the NE. A roughly cylindrical channel runs from top to bottom in the middle of each NPC (Stoffler et al., 2006). It is generally accepted that the central channel forms the pathway for the movement of large molecules, and it was assumed that the central channel would also allow the passage of small molecules and ions. However, many studies have shown that the binding of proteins that occlude the central channel and prevent active transport do not affect the passive diffusion of small molecules (Kramer et al., 2007). It has been demonstrated that there is an alternative pathway for the passage of ions and small molecular weight species through \sim eight symmetrical pores that are located around the periphery of NPCs (Shahin et al., 2001).

NPCs allow metabolites, nucleic acids and many proteins to pass efficiently across the NE via passive diffusion (Lee et al., 1998). Proteins in excess of ~ 50 kDa require energy-dependent movement across the NPC. Such large proteins are often designated for nuclear entry or exit by a specific NLS or NES. NLS and NES domains are constitutively displayed by some proteins to ensure continuous nuclear or cytosolic distribution. In other proteins, these targeting sequences can be revealed by covalent modification of the protein, thereby providing a mechanism for dynamic control of protein subcellular localisation. A classic example is the transcription factor NFAT (see main text for details) (Shaw, 1988; Hallhuber et al., 2006).

A commonly observed but controversial feature of NPCs is the presence of a 'central plug' that appears to occupy, and perhaps occlude, the channel (see figure). The nature of the putative NPC plug has been widely debated. Although it has been described as a dynamically regulated component of the NPC that somehow moves into place to occlude the pore, it has also been suggested that the plug is in fact cargo caught in transit. Support for the latter idea comes from the observation that there is an increased number of NPCs containing plugs when nuclei are cooled or if ATP is withdrawn, which prevents active transport (Stoffler et al., 1999). If the central plug were made up of randomly bound cargo, it could not be part of a deliberate Ca^{2+} -sensitive mechanism for regulating nuclear-cytosolic exchange of macromolecules. However, data that oppose the idea that the central plug is made up of randomly bound cargo include AFM studies showing that there are two reproducible conformations of the NPC. Given that different types of cargo probably have various shapes, AFM images of NPCs that had a greater variety of central plug conformations would be expected. However, it must be considered that most cargos are probably much smaller than the NPC, and therefore it is possible that they would not be evident in AFM studies.

Instead of using membrane-attached patches, Oberleithner and colleagues employed an 'hourglass' technique in which an isolated nucleus was sandwiched in a thin capillary tube. Using this method, it was observed that NPCs are constitutively open under physiological conditions (Danker et al., 2001). It was possible to modulate the NPC current recorded using the hourglass technique. For example, the addition of ATP or the depletion of the NE Ca^{2+} store increased the conductance of the NPC, but the NPCs always permitted cation flux (Shahin et al., 2001).

These latter findings arguably convey the physiological properties of NPCs most accurately. It is likely that patch-clamping of the NE distorts the membrane and somehow causes the NPCs to become non-conductive, whereas, under native conditions that are maintained using the hourglass technique, NPCs allow the constitutive passage of Ca^{2+} and other ions. Therefore, NPCs cannot be considered a complete diffusion break for Ca^{2+} . Rather, ions in the nucleoplasm or cytosol have permanent access to the other compartment (Eder and Bading, 2007). At most, NPCs might act as a diffusion filter and introduce a kinetic delay in the equilibration of nuclear-cytosolic Ca^{2+} concentrations. The extent of the kinetic delay might be subject to modulation: although NPCs do not close, their conductance can change in response to factors such as Ca^{2+} and ATP. Furthermore, the extent of NPC expression on the NE can vary from 1–50 NPCs per μm^2 , depending on cell type (Wang and Clapham, 1999). A greater expression of NPCs would allow a more rapid equilibration of cytosolic and nuclear Ca^{2+} concentrations.

Regulation of NPCs – a role for Ca^{2+} ?

Numerous studies have proposed that NPCs are regulated by changes in Ca^{2+} concentration either within the lumen of the NE, or at the cytosolic-nucleoplasmic face. However, there is little consistency in the nature of the observed effect. A widely cited example is the demonstration that altering the Ca^{2+} concentration in the lumen of the NE regulates both passive diffusion and NLS-mediated protein transport through NPCs (Greber and Gerace, 1995): depletion of the NE Ca^{2+} store was found to attenuate the nuclear influx of proteins bearing a NLS and of a non-specific fluorescent dextran molecule. It was suggested that the Ca^{2+} sensor in the NE was the integral membrane protein gp210, a nucleoporin of 210 kDa that functions to anchor NPCs. gp210 interacts with NPCs via its cytosolic C-terminus, whereas the bulk of the protein projects into the lumen of the NE, where it can sense Ca^{2+} levels and thereby mediate changes in NPC structure (Greber et al., 1990). In support of this idea, an antibody specific for the luminal domain of gp210 was found to inhibit both passive diffusion and signal-mediated transport into the nucleus (Greber and Gerace, 1992).

A similar finding was reported when the nuclear import of histone H1 (a ~21 kDa constitutively expressed nuclear protein) and two smaller fluorescently tagged dextran molecules (of 3 and 10 kDa) was compared in intact cardiomyocytes (Perez-Terzic et al., 1999). Reducing the Ca^{2+} concentration in the NE lumen by treating cells with a Ca^{2+} chelator, a Ca^{2+} -pump inhibitor or ionomycin prevented the energy-dependent import of the histone. The same treatments prevented the passive influx of 10 kDa dextran, but not the 3 kDa dextran species. These data indicate that NPC function is sensitive to the concentration of Ca^{2+} within the NE, and that altering luminal Ca^{2+} concentration can regulate both active and passive flux of large, but not small, molecules. However, a similar study examining the diffusion of GFP between the nucleoplasm and cytosol of intact *Xenopus* oocytes reached the opposite conclusion; the NE reduced

diffusion of GFP by ~100-fold, but depletion of Ca^{2+} from the lumen of the NE made no difference (Wei et al., 2003). Yet a different finding was reported in a study of the movement of photo-activated GFP across the NE in hepatocytes (O'Brien et al., 2007). In this case, a rise in cytosolic Ca^{2+} levels, and not changes in NE Ca^{2+} content, increased the influx (but not the efflux) of photo-activated GFP across the NE. In this latter study, it was also demonstrated that Ca^{2+} signals evoked by InsP_3 Rs regulated the influx of GFP into the nucleus, suggesting a putative link between cell activation by InsP_3 -producing stimuli and the control of nuclear trafficking.

These discrepant studies highlight some of the controversy surrounding how NPC opening is regulated. As yet, there is little consensus about what effects cytosolic Ca^{2+} , or Ca^{2+} within the NE, have on NPC structure. However, although the results of the various studies are not entirely consistent, they suggest that Ca^{2+} is somehow important in regulating nuclear traffic. The mechanism by which Ca^{2+} might regulate NPC permeability is unclear. One of the most commonly proposed theories is that a central 'plug' moves and blocks the channel in response to changes in the Ca^{2+} concentration in either the cytosol or the lumen of the NE (see Box 3). Consistent with this idea, several studies reported Ca^{2+} -dependent changes in the shape of NPCs (Stoffler et al., 1999) and, in particular, alterations in the number of NPCs bearing a central plug. A key technique in these investigations was atomic force microscopy (AFM), which can be used to scan the three-dimensional topography of the cytosolic face of NPCs. AFM can resolve changes in the surface of large macromolecules, and has been used to show that the depletion of Ca^{2+} from the lumen of the NE significantly and reversibly increased the proportion of NPCs with a central plug (Wang and Clapham, 1999). The observed change in NPC structure that is caused by luminal Ca^{2+} depletion is in line with the Ca^{2+} -dependent alterations in NPC transport described above (Perez-Terzic et al., 1999), and supports the concept of allosteric regulation of NPC gating by Ca^{2+} . However, the reported changes in NPC structure are not the same in different studies (Erickson et al., 2006), and alteration in NPC structure following Ca^{2+} release from the NE is not universally observed (Stoffler et al., 2006).

Even if changes in NPC permeability do indeed occur under physiological conditions, it is not clear what difference this makes to cellular growth or activity. It has been argued that simply altering the permeability of the nucleus to species of different molecular masses is a rather crude mechanism for regulating nuclear-to-cytosolic flux (Torok, 2007). Much greater specificity for NPC transit can be achieved by cellular signalling mechanisms that either promote or diminish the exposure of molecular tags (such as the NLS and NES) that promote NPC transport.

Nuclear Ca^{2+} signalling in intact cells

It has been established that Ca^{2+} signals occur in the nucleus of intact cells. For example, confocal imaging studies have shown that as Ca^{2+} waves 'sweep' through the cytosol of hormone-stimulated cells, the nucleus becomes engulfed by the signal (Lipp et al., 1997). However, a major controversial issue concerning nuclear Ca^{2+} signalling is whether Ca^{2+} transients are generated autonomously in the nucleoplasm, and whether they can be distinct from cytosolic Ca^{2+} changes. Numerous studies have examined Ca^{2+} signalling in isolated nuclei (Adebanjo et al., 2000; Bezin et al., 2008b; Gerasimenko et al., 1995). The advantage of studying nuclei in isolation is that changes in nucleoplasmic Ca^{2+} only occur if Ca^{2+} is released from channels within the inner NE. Although the NPCs are present in this type of preparation, it is unlikely that Ca^{2+} released

from channels on the outer NE (that face away from the nucleus), or from potential remnants of the ER, can diffuse into the nucleoplasm. Rather, any Ca^{2+} that is released outside of the NE is rapidly dissipated and diluted. Isolated nuclei therefore represent a system in which nucleoplasm-directed Ca^{2+} -release channels can be examined. However, although these studies are useful for examining the potential of nuclei to generate their own Ca^{2+} signals, they do not determine whether these signals are at all significant compared with the larger fluxes of Ca^{2+} that occur in the cytosol of intact cells. The key issue, therefore, is to establish how nuclear Ca^{2+} signals arise within the physiological context of intact cells, and whether Ca^{2+} channels facing the nucleoplasm have a significant role.

Measuring nuclear Ca^{2+} signals: key considerations

The idea that nuclear Ca^{2+} levels are regulated independently from those in the cytosol emerged as soon as imaging technology advanced to allow monitoring of Ca^{2+} changes simultaneously in different cellular compartments (Bkaily et al., 1997). Using both wide-field and confocal fluorescence imaging of living cells, several groups began to report that nuclear Ca^{2+} concentration differed from that in the cytosol. The extent to which the Ca^{2+} signals in these two compartments differed was highly variable between studies. In some cases, the nuclear and cytosolic Ca^{2+} levels were found to be entirely independent, whereas others reported modest differences in the kinetics or amplitude of the nucleoplasmic and cytosolic responses (Leite et al., 2003; Santella et al., 2003). Although the examples of differential Ca^{2+} concentration in the nucleus and cytosol are too numerous to mention in detail, some are discussed below, and Table 1 lists some of the reported findings.

Owing to their brightness, Ca^{2+} -sensitive fluorescent indicators have been used in most studies that compared the properties of nuclear and cytosolic Ca^{2+} signals. However, it is certain that many of these studies are invalid as they failed to recognise that organic fluorochromes have different characteristics in the nuclear and cytosolic compartments (Birch et al., 1992; Bkaily et al., 2006). Indeed, one of the most prominent early studies of nuclear Ca^{2+} regulation – which suggested that Ca^{2+} concentration in the nucleus tracked that in the cytosol up to ~300 nM, after which they were insulated against further rises (al-Mohanna et al., 1994) – is compromised by the lack of in situ calibration (O'Malley et al., 1999). We (Thomas et al., 2000b) and others (O'Malley et al., 1999; Perez-Terzic et al., 1997) have demonstrated that commonly used fluorescent Ca^{2+} indicators are much brighter in the nucleus than in the cytosol; they differ with respect to both their affinity for Ca^{2+}

and their dynamic range. Therefore, to obtain absolute values for Ca^{2+} concentration in different cellular compartments, it is crucial to perform independent calibrations with the indicator in exactly the same environment in which it will be used (Thomas et al., 2000b). Typically, when quiescent cells are loaded with a Ca^{2+} indicator such as fura2 or fluo3, it is evident that the nucleus has a greater fluorescence than the cytosol. The difference in the intensity of the indicator fluorescence between the nucleus and cytosol is maintained or even exaggerated as the Ca^{2+} concentration globally increases. This exaggerated nuclear fluorescence has led some studies to erroneously conclude that nuclei amplify cellular Ca^{2+} signals, and even that there are intranuclear Ca^{2+} gradients (Birch et al., 1992). It is not known why fluorescent Ca^{2+} indicators behave differently in the nucleus, although it has been established that factors such as ionic strength, viscosity and pH can alter Ca^{2+} binding to the indicator. In addition, it appears that Ca^{2+} indicators interact with nuclear contents; for example, in *Xenopus* oocyte nuclei that had been loaded with fluo3, it was found that the indicator did not diffuse freely because it became somehow 'trapped' in chromatin and/or DNA (Perez-Terzic et al., 1997). These technical artefacts are one of the main reasons why nuclear Ca^{2+} signalling has such a controversial history.

Perinuclear Ca^{2+} release can affect nucleoplasmic Ca^{2+} levels

The observation of a nucleoplasmic Ca^{2+} signal does not necessarily indicate that the nuclei themselves are the actual source of Ca^{2+} . It is evident that Ca^{2+} diffuses more slowly in the cytosol than in the nucleoplasm because of the much greater buffering and sequestration capacity in the cytosolic compartment (Fox et al., 1997). This means that Ca^{2+} signals that arise immediately outside of the nucleus have a greater potential to diffuse through NPCs and into the nucleus than to permeate across the cytosol. Consistent with this idea, it was observed that mobilisation of Ca^{2+} from intracellular stores surrounding the nucleus of various cell types caused simultaneous Ca^{2+} elevations within the cytosol and nucleoplasm (Chamero et al., 2002), whereas distal Ca^{2+} signals that arose from voltage-operated Ca^{2+} channels at the plasma membrane induced cytosolic Ca^{2+} transients, but only modest and delayed increases of nucleoplasmic Ca^{2+} .

An emerging concept in Ca^{2+} signalling is that localised perinuclear Ca^{2+} release – that is, microscopic Ca^{2+} release events that occur next to the nucleus – can affect nuclear activities (Fig. 3A). It has been shown that brief openings of InsP_3Rs or RyRs induce microscopic Ca^{2+} events that do not diffuse farther than a

Table 1. Selected examples of observed differences in cytosolic and nucleoplasmic Ca^{2+} levels

Nature of difference in nuclear and cytosolic Ca^{2+} signals	Cell type	Experimental method	References
Temporally independent Ca^{2+} signals occurred in each compartment	Smooth muscle	Fluorescence imaging	(Fedoryak et al., 2004)
Lower threshold for InsP_3 -evoked Ca^{2+} release in the nucleus	PC12, NIH3T3 cells	Fluorescence imaging	(Huh et al., 2007)
Nucleoplasmic Ca^{2+} signal lagged ~4 seconds behind elevation in cytosolic Ca^{2+}	Starfish oocytes	Fluorescence imaging	(Santella et al., 2003)
Nucleoplasmic Ca^{2+} signal lagged ~100 milliseconds behind elevation in cytosolic Ca^{2+}	Cardiac myocytes	Fluorescence imaging	(Bootman et al., 2007; Kockskamper et al., 2008a)
Lower threshold for InsP_3 -evoked Ca^{2+} release in the nucleus	Liver-cell line	Fluorescence imaging	(Leite et al., 2003; Echevarria et al., 2003)
Larger increase in Ca^{2+} in the nucleoplasm than in the cytosol, but with the same kinetics	HeLa cells	Bioluminescence (aequorin)	(Brini et al., 1993)
Smaller increase in Ca^{2+} in nucleoplasm than in the cytosol, but with the same kinetics	HeLa cells	Bioluminescence (aequorin)	(Badminton et al., 1996; Badminton et al., 1995)
Nucleoplasmic Ca^{2+} signals occurred in the absence of Ca^{2+} changes in the surrounding cytosol	Cardiac myocytes	Fluorescence imaging	(Higazi et al., 2009)




Fig. 3. Perinuclear Ca^{2+} channels can affect nucleoplasmic Ca^{2+} concentration. (A) The differential effect of perinuclear versus distal Ca^{2+} release on nucleoplasmic Ca^{2+} levels. When Ca^{2+} puffs are activated distal to the nucleus, they are constrained to a distance of $\sim 2\text{--}4\ \mu\text{m}$ owing to cytosolic Ca^{2+} buffering and sequestration. These Ca^{2+} puffs do not diffuse through the cytosol, and therefore do not reach the nucleus. By contrast, when a Ca^{2+} puff of similar amplitude occurs next to the nucleus, it diffuses anisotropically into the nucleoplasm, engulfs the entire nucleus and can even spill out of the nucleus on the opposite side to where the Ca^{2+} signal originated. Further details can be found in (Lipp et al., 1997). (B) Perinuclear expression of type 2 InsP_3Rs in a neonatal myocyte.

few micrometres. If such events occur in the cytosol at sites that are distant from the nucleus, they probably have only a modest impact on cell behaviour, but if they occur next to the nucleus, then they can have a long-lasting and integrating effect on nucleoplasmic Ca^{2+} . Such observations have been made using both excitable and non-excitable cell types (Lipp et al., 1997). Indeed, many of the studies that proposed the existence of independent nuclear Ca^{2+} signals could be explained by such a perinuclear cytosolic Ca^{2+} release, followed by anisotropic diffusion of Ca^{2+} through NPCs into the nucleoplasm. In particular, this probably applies to studies in which the nuclear Ca^{2+} signal lags behind the cytosolic Ca^{2+} response (Huh et al., 2007).

Perinuclear Ca^{2+} release, and the consequent elevation in nuclear Ca^{2+} that it induces, has recently become a particular focus in studies of cardiac myocytes (Higazi et al., 2009; Kocksamper et al., 2008b; Wu et al., 2006). Ca^{2+} signals occur rhythmically in cardiac myocytes and promote the interaction of protein filaments that make the cells contract and allow the heart to pump blood. This process of excitation-contraction (EC) coupling in the heart relies on the interplay between voltage-gated Ca^{2+} channels on the plasma membrane and RyRs on the sarcoplasmic reticulum (the major internal Ca^{2+} store). The voltage-gated Ca^{2+} channels respond to the depolarizing action potential that sweeps over the heart with each beat by opening and thereby permitting the influx of a small amount of Ca^{2+} . This Ca^{2+} signal is greatly amplified by RyRs that

are close to the voltage-gated Ca^{2+} channels. Ca^{2+} diffuses from the RyRs to the actin/myosin filaments and initiates cell contraction. RyRs are therefore considered to be the main Ca^{2+} -release channel in the heart. However, cardiac myocytes also express InsP_3Rs (Lipp et al., 2000; Perez et al., 1997), and recent studies of neonatal (Garcia et al., 2004; Guatimosim et al., 2008; Luo et al., 2008b; Luo et al., 2006), atrial (Bootman et al., 2007; Kocksamper et al., 2008a; Zima et al., 2007) and ventricular myocytes (Higazi et al., 2009; Wu et al., 2006) demonstrated that InsP_3 -generating agonists, or InsP_3 itself, have specific effects on nucleoplasmic Ca^{2+} signals. Indeed, it appears that a substantial proportion of InsP_3Rs in neonatal and adult cardiomyocytes is expressed on membranes close to the nucleus or on the NE (Bare et al., 2005; Higazi et al., 2009; Liu et al., 2001) (Fig. 3B). This strategic positioning of InsP_3Rs might allow the generation of nucleus-specific Ca^{2+} signals (even though they originate in the cytosolic compartment), which could have a significant role in regulating cardiac gene transcription, and thereby be involved in controlling processes such as cardiac hypertrophy. In support of this, it has been demonstrated that the release of perinuclear InsP_3 -sensitive Ca^{2+} stores promotes the activation of calcineurin and nuclear transport of NFAT into the nucleus to trigger a hypertrophic gene programme (Higazi et al., 2009). In addition, perinuclear InsP_3 -mediated Ca^{2+} release promotes the phosphorylation of histone deacetylase 5 via Ca^{2+} -calmodulin-dependent kinase II, causing it to be exported out of the nucleus (Wu et al., 2006) and thereby de-repressing genes that underlie the hypertrophic growth of myocytes.

It is well known that the stimulation of myocytes with hormones that activate InsP_3 production can promote hypertrophy (Sugden and Clerk, 2005), but it is not clear how hormones induce subtle changes in Ca^{2+} -dependent gene transcription within cells that already experience periodic surges of Ca^{2+} during EC-coupling (Roderick et al., 2007). Perinuclear InsP_3Rs might explain how this occurs, as they can provide a source of Ca^{2+} that is spatially and temporally distinct from the Ca^{2+} signals that are involved in EC-coupling (Molkentin, 2006). Consistent with this model, stimulation of rat neonatal myocytes with phenylephrine evoked InsP_3R -mediated perinuclear Ca^{2+} -release events that entered the nucleoplasm (Luo et al., 2008). Together with the finding that inhibiting InsP_3R opening prevented hypertrophic growth in response to phenylephrine stimulation (Luo et al., 2006), it is logical to conclude that phenylephrine promotes hypertrophy by activating perinuclear InsP_3Rs and, consequently, causing nucleoplasmic Ca^{2+} signals. Comparable observations have been made in cardiac cells stimulated with another InsP_3 -generating agonist, endothelin-1; InsP_3Rs in close proximity to the NE triggered nucleoplasmic Ca^{2+} signals and hypertrophic gene transcription, whereas EC-coupling did not (Garcia et al., 2004; Higazi et al., 2009). Although InsP_3Rs conduct significantly less Ca^{2+} and are expressed at levels that are ~ 100 -fold lower than RyRs, the activation of InsP_3Rs can clearly have a dramatic effect on gene transcription in cardiac myocytes owing to their specific impact on nuclear Ca^{2+} . It remains to be determined whether this model of perinuclear Ca^{2+} release is involved in regulating gene transcription and other nuclear activities in other cell types. However, there are numerous studies that have examined InsP_3R distribution in various cell types and shown that these Ca^{2+} channels are often most densely concentrated around the nucleus (Bootman et al., 2001; Shirakawa and Miyazaki, 1996; Thomas et al., 2000a; Vermassen et al., 2003). It therefore seems probable that modulation of nucleoplasmic Ca^{2+} concentration by perinuclear Ca^{2+} release is a generic signalling paradigm.

Independent nuclear Ca^{2+} signalling

The studies described above substantiate the idea that perinuclear Ca^{2+} release significantly affects nucleoplasmic Ca^{2+} concentration. However, the observations also raise questions about the function of Ca^{2+} channels located on the inner NE and in the nucleoplasmic reticulum. In particular, why don't the Ca^{2+} channels on the inner NE, or other putative nuclear Ca^{2+} stores, generate a Ca^{2+} signal that is independent from the cytosol? Or do Ca^{2+} signals that arise in the cytosol always dominate what happens in the nucleus? As mentioned earlier, it has been shown that isolated nuclei have the machinery to release Ca^{2+} , but the majority of intact cell studies indicate that nucleoplasmic Ca^{2+} signals originate outside the nucleus (Lipp et al., 1997; Luo et al., 2008). Indeed, in one notable study, InsP_3 was introduced specifically to the nucleus or cytosol to examine the relative ability of this messenger to trigger Ca^{2+} signals in each compartment. As expected, when released in the cytosol, InsP_3 evoked a cytosolic Ca^{2+} signal that invaded the nucleus via diffusion. However, when InsP_3 was released in the nucleus, the resulting Ca^{2+} signal also initiated in the cytosol, and then invaded the nucleus (Shirakawa and Miyazaki, 1996). This indicates that even when InsP_3 is given immediate access to the InsP_3 Rs on the NE, the messenger diffuses out of the nucleus and preferentially triggers cytosolic Ca^{2+} release. A similar conclusion was reached when an InsP_3 R antagonist was introduced in the cytosol of cells during stimulation with an InsP_3 -generating agonist. The antagonist, which was too large to enter the nucleus, blocked both cytosolic and nucleoplasmic Ca^{2+} transients (Allbritton et al., 1994). The obvious conclusion from such studies is that, although nuclei have the capacity to generate Ca^{2+} transients, their contribution is 'swamped' by the larger cytosolic Ca^{2+} release [but see Lui et al. (Lui et al., 1998) for a contradictory observation].

Perhaps it is only under specific stimulation conditions that substantial Ca^{2+} responses are evoked from nuclear InsP_3 Rs. For example, a comparison between the Ca^{2+} signals that are evoked by vasopressin and hepatocyte growth factor (HGF) in hepatic carcinoma cells revealed that vasopressin triggered Ca^{2+} release in the cytosol, whereas HGF caused nuclear Ca^{2+} mobilisation (Gomes et al., 2008). In the latter case, it was suggested that the HGF receptor was internalised and translocated to the nucleus, where it triggered local InsP_3 production. Vasopressin, on the other hand, stimulated InsP_3 production in the cytosol. It is therefore plausible that different stimuli activate InsP_3 Rs in alternative cellular compartments, depending on where in the cell the InsP_3 is generated.

An example of nuclear-specific Ca^{2+} signalling is presented in Fig. 4A, which shows the spatial profile of spontaneous Ca^{2+} transients that occur in a neonatal cardiac myocyte. Such Ca^{2+} signals are largely generated by the stochastic activation of InsP_3 Rs and RyRs (Luo et al., 2008). The black trace shows substantial nuclear Ca^{2+} elevations, but a corresponding increase in cytosolic Ca^{2+} on either side of the nucleus is not observed. It was not determined whether the nuclear Ca^{2+} signals were due to activation of Ca^{2+} channels on the inner NE, or to perinuclear Ca^{2+} release. However, the figure presents an example in which substantial Ca^{2+} changes can be clearly observed solely in the nucleus.

Autonomous generation of Ca^{2+} -mobilising messengers in the nucleus

If nuclei are truly independent of cytosolic signalling, they must have the ability to generate Ca^{2+} -mobilising messengers that then act locally on nucleoplasmic InsP_3 Rs, RyRs and NAADPRs.




Fig. 4. Cytosolic and nuclear Ca^{2+} signals can be independent. (A) An example of Ca^{2+} signals that occur in the nucleus of a neonatal rat cardiac myocyte independently of Ca^{2+} signals in the cytosol. In the inset, two adjacent cells are outlined in white. The cells showed spontaneous Ca^{2+} -release events, which is typical of neonatal cardiomyocytes in culture. The Ca^{2+} signals were recorded by loading the cells with fluo4 and monitoring changes in fluorescence emission using confocal microscopy, and the responses were recorded from the positions indicated in the left-hand cell. Calcium increase is shown as the fluo4 fluorescence normalised to the baseline (F/F_0). (B) Difference in the kinetics of cytosolic and nuclear Ca^{2+} in an adult atrial cardiac myocyte during regular electrical pacing. Cells were paced to steady state before the recording was started. The traces were obtained by confocal line-scanning of fluo4 fluorescence across a region perpendicular to the long axis of the cell. Line scans were obtained at 2-ms intervals using an Olympus FluoView 1000. The regions of the line scan from which the cytosolic and nuclear Ca^{2+} traces in C were obtained are indicated on the left of the line scan image (marked by 'C'). (C) Spatial properties of the cytosolic and nuclear Ca^{2+} signals. The intensity of the white pixels is proportional to Ca^{2+} concentration, as shown in the right-hand panel.

Numerous studies have demonstrated that the biochemical machinery that is required for the generation of Ca^{2+} -mobilising messengers is present in the nucleus. For example, the ADP-ribosyl cyclase enzyme that is responsible for the production of cyclic ADP ribose (cADPR) and/or NAADP is localised in the nucleus (Adebanjo et al., 1999; Bezin et al., 2008a; Trubiani et al., 2008). In addition, it has been well established that nuclei possess phosphoinositide signalling mechanisms that are similar to, but distinct from, those that occur at the plasma membrane, and which can lead to InsP_3 production (Visnjic and Banfic, 2007; Ye and Ahn, 2008). Indeed, nuclei have substantial amounts of the membrane phospholipid phosphatidylinositol 4,5-bisphosphate [$\text{PtdIns}(4,5)\text{P}_2$]

and phosphoinositide-specific phospholipase C (PtdIns-PLC) (Downes et al., 2005). Interestingly, it has been reported that PtdIns(4,5) P_2 and PtdIns-PLC localise in particles known as nuclear speckles (see Fig. 1). These small nuclear substructures are believed to be the sites of storage and/or assembly of pre-mRNA splicing factors. The exact function of PtdIns(4,5) P_2 and PtdIns-PLC in nuclear speckles is currently under investigation, but they are involved in linking the production of diacylglycerol (DAG) and $InsP_3$ (and their metabolites) to the control of RNA processing (Alcazar-Roman and Wente, 2008).

Mammalian cells express at least 13 different PtdIns-PLC isozymes that can be grouped into six subfamilies (β , γ , δ , ϵ , η and ζ) (Cockcroft, 2006; Suh et al., 2008), of which PtdIns-PLC β_1 is considered to be the predominant nuclear form (Faenza et al., 2008; Ye and Ahn, 2008). In some cell types, PtdIns-PLC β_1 is constitutively present in nucleus, whereas in others, growth-factor stimulation causes the enzyme to translocate from the cytosol to the nucleus. This is facilitated by C-terminal sequences in PtdIns-PLC β_1 that act either as a NLS or cause the retention of the protein in the nucleus (Kim et al., 1996). It is known that PtdIns-PLC β isoforms are activated at the plasma membrane by G-protein-coupled receptors and are responsible for rapid $InsP_3$ production and acute Ca^{2+} signals (see Box 2). In the nucleus, however, PtdIns-PLC β_1 plays an important role in signalling events that are evoked by growth-factor-receptor stimulation (Visnjic and Banfic, 2007). For example, PtdIns-PLC β_1 was found to be essential during the differentiation of skeletal muscle myoblasts (Ramazzotti et al., 2008). The expression of the enzyme increases during myogenesis, and is required for the induction of cyclin D3 expression and the formation of the retinoblastoma-cyclin D3 complex that underlies differentiation (De Santa et al., 2007). The nuclear localisation of PtdIns-PLC β_1 is essential in this context, as a cytosolically targeted form does not support myoblast differentiation (Faenza et al., 2007).

These studies highlight the importance of nuclear PtdIns-PLCs in growth factor responses. However, they mainly describe the roles of nuclear DAG, protein kinase C translocation or inositol polyphosphate formation that results from PtdIns-PLC activation (Divecha et al., 1991). Considerably less attention has been focussed on whether changes in the levels of nuclear Ca^{2+} participate in growth-factor-mediated responses. However, several recent reports have suggested a crucial role for Ca^{2+} downstream of nuclear PtdIns-PLC activation. For example, insulin receptors are found in the nucleus following insulin stimulation of cells, where they promote PtdIns(4,5) P_2 turnover and nuclear Ca^{2+} signalling independently of cytosolic Ca^{2+} signalling (Rodrigues et al., 2008). The specific action of insulin on nuclear $InsP_3$ production was confirmed by expressing the $InsP_3$ -binding portion of the $InsP_3$ R (known as an $InsP_3$ 'sponge') to buffer changes in $InsP_3$ concentration. To affect its subcellular location, the $InsP_3$ sponge was fused to either a NLS or a NES. The expression of the nucleus-targeted $InsP_3$ sponge significantly attenuated responses to insulin, whereas the cytosol-targeted $InsP_3$ sponge did not (Rodrigues et al., 2008). The opposite effect was observed if the cells were stimulated with vasopressin, which acts on the plasma membrane pool of PtdIns(4,5) P_2 via cytosolic PtdIns-PLC isozymes (Gomes et al., 2008).

In addition to growth factor receptors being present in the nucleus, a number of studies have demonstrated that functional hormone receptors are also internalised and move across the NE. For example, a C-terminal NLS directs the nuclear import of the bradykinin B2 receptor (Savard et al., 2008). Specific bradykinin-binding sites are present in hepatocyte nuclear extracts, and the

addition of bradykinin to isolated nuclei was found to evoke a rapid, transient nucleoplasmic Ca^{2+} signal. Furthermore, it has been demonstrated that endogenous and ectopically expressed mGluR5 metabotropic glutamate receptors are present on the nuclear membranes of primary neurons and of cultured cells, respectively (Kumar et al., 2008). Studies of isolated nuclei indicate that these receptors activate PtdIns-PLCs in the nucleus, leading to nucleoplasmic $InsP_3$ generation and Ca^{2+} signals.

Exactly how G-protein-coupled receptors in the nucleus become activated is unclear. The lumen of the NE is topologically the same as the extracellular space, so the ligand-binding domain of nuclear hormone receptors would be exposed within the NE lumen. It has been proposed that specific transporters allow the uptake of neurotransmitters such as glutamate into the NE, where they can bind to their cognate receptors and activate signalling (Jong et al., 2005). Similarly, receptors for endothelin, angiotensin and prostaglandin, the activation of which can also stimulate Ca^{2+} -mobilising pathways, are reportedly expressed on nuclear membranes (Bkaily et al., 2006; Coffey et al., 1997).

Although many studies have focussed on investigating the nuclear functions of PtdIns-PLC β_1 , other PtdIns-PLC isoforms are also found in the nucleus, in particular PLC δ_1 (Okada et al., 2005) and PtdIns-PLC γ_1 (Gomes et al., 2008). In quiescent cells, PtdIns-PLC δ_1 is localised at the plasma membrane and in the cytosol, but following stimulation, it translocates to the nucleus in a Ca^{2+} -dependent manner (Okada et al., 2005), where it plays a role in regulating PtdIns(4,5) P_2 levels, DNA synthesis and cell proliferation (Stallings et al., 2008). Growth-factor-induced Ca^{2+} signals, as well as non-specific Ca^{2+} transients caused by ionophores, promote nuclear PtdIns-PLC δ_1 translocation, thereby acting as positive feedback for further nuclear Ca^{2+} signalling (Yagisawa, 2006). Although data regarding the functions of different PtdIns-PLC isozymes in the nucleus are still emerging, current evidence suggests that they all play roles in cell survival, division and differentiation.

PtdIns-PLC ζ is also known to accumulate in the nucleus owing to an NLS, but in this case, the translocation of the enzyme prevents Ca^{2+} signalling (Ito et al., 2008). Repetitive Ca^{2+} oscillations occur immediately following fertilisation in mammalian oocytes, and are believed to be driven by $InsP_3$ production that is caused by the introduction of PtdIns-PLC ζ into the oocyte from the fused sperm. The Ca^{2+} oscillations cease around the time when the pronucleus is formed, owing to the sequestration of PtdIns-PLC ζ in the nucleus (Larman et al., 2004). Unlike the other PtdIns-PLC isozymes described above, when PtdIns-PLC ζ is in the nucleus, it does not evoke $InsP_3$ production and Ca^{2+} signalling.

The Ca^{2+} -mobilising pathways described above have well-characterised effects on nuclear Ca^{2+} signalling. However, there are numerous additional messengers and signalling pathways by which cellular Ca^{2+} signals can be evoked (Bootman et al., 2002), and some of these have been shown to function in the nucleus. Another example of a putative nuclear messenger is arachidonic acid: this lipid is known to increase cytosolic Ca^{2+} concentration. Although little is known about the effect of arachidonic acid on nucleoplasmic Ca^{2+} , it is plausibly an additional regulator of nuclear Ca^{2+} concentration. Indeed, phospholipase A2, an enzyme involved in the production of arachidonic acid, translocates into the nucleus in a Ca^{2+} -dependent manner (Schievella et al., 1995). Furthermore, arachidonic acid can be metabolised into further Ca^{2+} signalling moieties by the action of cyclooxygenase-2, which translocates into the nucleus in response to growth-factor-stimulation (Coffey et al., 1997). Although the bulk of studies have focussed on $InsP_3$ -mediated Ca^{2+} release in

the nucleus, it is probable that nuclear Ca^{2+} signalling utilises the same diverse range of messengers and channels as the cytosol.

Recovery of nucleoplasmic Ca^{2+} signals

It has been noted in several studies that nuclear Ca^{2+} signals persist considerably longer than equivalent cytosolic Ca^{2+} transients (Bootman et al., 2007; Kockskamper et al., 2008a; Lipp et al., 1997). The reason for this is the lack of Ca^{2+} ATPases on the inner NE (Humbert et al., 1996). Because of the slow dissipation of nuclear Ca^{2+} , it is plausible that rapid cellular stimulation could cause a progressive accumulation of Ca^{2+} in the nucleoplasm. Fig. 4B,C shows an example of the kinetic delay in both the rise and dissipation of nuclear Ca^{2+} . The recordings were obtained by imaging fluo4 fluorescence from a single atrial cardiac myocyte that was activated by regular electrical depolarisation. In addition to the kinetic delay in recovery of the nuclear Ca^{2+} transient, it is evident that the nuclear fluo4 fluorescence was consistently higher than that in the cytosol, even when both signals had recovered. This is because fluo4 is brighter in the nucleus than in the cytosol (as described above), not because there was a persistent gradient of Ca^{2+} between the cytosol and nucleus. The calibrated diastolic Ca^{2+} levels are actually very similar in the cytosol and nucleus. The differences in the kinetics of the nuclear and cytosolic Ca^{2+} signals are illustrated in the line-scan image shown in Fig. 4C; it is evident that the nucleus takes longer to fill with Ca^{2+} , and considerably longer for the Ca^{2+} signal to dissipate.

A major mechanism for the dissipation of nuclear Ca^{2+} signals is via simple diffusion through NPCs. To leave the nucleus, a Ca^{2+} ion has to pass through the nucleoplasm and then encounter and traverse a NPC. Once outside the nucleus, Ca^{2+} can be quickly sequestered by Ca^{2+} ATPases on the outer NE, ER and plasma membrane, or by other buffers such as mitochondria. The NE can be regarded as a diffusion barrier for both Ca^{2+} entry and exit. Although Ca^{2+} ATPases are not expressed on the inner NE, there is growing evidence that a splice variant of type 1 sodium-calcium exchanger (NCX) is expressed on the inner NE (Ledeen and Wu, 2007). It is suggested that the nuclear NCX uniquely associates with a ganglioside known as GM1, which potentiates Ca^{2+} transport and prevents Ca^{2+} overloading within the nucleus (Ledeen and Wu, 2008). NCX is a mechanism of Ca^{2+} transport that is usually associated with electrically excitable cells in which large and rapid fluxes of Ca^{2+} occur. However, many non-excitable cells also express NCX and, in some cells, expression of the transporters appears to be restricted to the nucleus (Xie et al., 2004). The configuration of the NCX on the inner NE is the same as on the plasma membrane, but the interaction with GM1 is atypical. In the presence of a Na^+ gradient, nuclei sequester Ca^{2+} , verifying the functionality of NCX (Xie et al., 2002). The Na^+ gradient might be driven by a Na^+/K^+ ATPase or Na^+/H^+ exchanger, all of which are also present on the NE (Bkaily et al., 2006; Garner, 2002). Therefore, in addition to passive diffusion out of the nucleus, nucleoplasmic Ca^{2+} signals can be reversed by active uptake processes.

Replenishment of Ca^{2+} in the NE lumen is probably mediated by ER-localised Ca^{2+} ATPases. Although these enzymes sequester cytosolic Ca^{2+} into the ER, the fact that the lumen of the NE and ER are connected means that Ca^{2+} can diffuse from one membrane system to the other (Wu and Bers, 2006). In addition, it has been reported that Ca^{2+} accumulates in the NE via a curious mechanism that is driven solely by InsP_4 (Hsu et al., 1998; Malviya and Klein, 2006). It has been reported that specific InsP_4 receptors are present on the outer NE (Malviya, 1994), but exactly how these binding sites activate Ca^{2+} uptake is unknown. A major problem with this

proposed uptake pathway is that the ER and NE possess a substantially higher Ca^{2+} concentration than the cytosol, and, unless the Ca^{2+} content of these stores was completely emptied, the flux of Ca^{2+} would always be towards the cytosol. Complete depletion of ER Ca^{2+} is a pathological state that culminates in cell death.

In the process of signal transduction, the reversal of a signalling event is as important as its induction. Exactly how nucleoplasmic Ca^{2+} signals decline back to pre-stimulation levels is not entirely clear, as the majority of studies have focussed solely on the generation of nucleoplasmic Ca^{2+} transients. Currently, passive diffusion through NPCs and uptake into the NE by NCX appear to be the main mechanisms by which nuclear Ca^{2+} transients are limited and reversed.

Conclusions

Over the past decade there has been a tremendous increase in the knowledge of Ca^{2+} signalling systems in the nucleus, and progress has been made in unravelling the impact of changes in nuclear Ca^{2+} on cellular activity. Recent studies have clearly highlighted the idea that gene transcription and cell-cycle progression can be specifically influenced by nuclear Ca^{2+} signals. It is probable that many more cellular processes will be shown to be dependent on nucleoplasmic Ca^{2+} changes. Despite this understanding of the functions of nuclear Ca^{2+} signals, there is still some uncertainty concerning how nuclear Ca^{2+} signals are generated and whether Ca^{2+} signalling in the nucleus is really autonomous. This is a surprisingly contentious issue that has been debated for many years. It has been shown that NPCs do not prevent Ca^{2+} fluxes from crossing the NE, which would imply that Ca^{2+} signalling in the nucleus must follow cytosolic Ca^{2+} changes (providing that cytosolic Ca^{2+} changes persist for sufficient time). Yet, several studies have shown that Ca^{2+} signals initiate within, and are restricted to, the nucleus. Exactly how Ca^{2+} signals within the nucleus or cytosol are prevented from equilibrating with the other compartment remains unclear.

In future studies, it will be important to work towards defining unified views on several topics, including nuclear Ca^{2+} -release mechanisms and the regulation of NPC structure and function by Ca^{2+} . As some of the controversy in nuclear Ca^{2+} signalling might be owing to the use of different cell types in different laboratories, it will also be necessary to consider cell-type-specific versus generic Ca^{2+} signalling paradigms. Finally, another goal of future research will be to link putative nucleoplasmic Ca^{2+} signals with cellular processes.

The authors would like to thank Simon Walker and Hattie O'Nions for their help in the preparation of Fig. 2 and Daniel Higazi for preparation of Fig. 3B. This work was supported by the BBSRC and the British Heart Foundation (grants PG/06/034/20637 and PG/07/040). H.L.R. is a Royal Society University Research Fellow. I.S. is a BBSRC-funded PhD student.

References

- Adebanjo, O. A., Anandatheerthavarada, H. K., Koval, A. P., Moonga, B. S., Biswas, G., Sun, L., Sodam, B. R., Bevis, P. J., Huang, C. L., Epstein, S. et al. (1999). A new function for CD38/ADP-ribosyl cyclase in nuclear Ca^{2+} homeostasis. *Nat. Cell Biol.* **1**, 409–414.
- Adebanjo, O. A., Biswas, G., Moonga, B. S., Anandatheerthavarada, H. K., Sun, L., Bevis, P. J., Sodam, B. R., Lai, F. A., Avadhani, N. G. and Zaidi, M. (2000). Novel biochemical and functional insights into nuclear Ca^{2+} transport through IP(3)Rs and RyRs in osteoblasts. *Am. J. Physiol. Renal Physiol.* **278**, F784–F791.
- al-Mohanna, F. A., Caddy, K. W. and Bolsover, S. R. (1994). The nucleus is insulated from large cytosolic calcium ion changes. *Nature* **367**, 745–750.
- Alcazar-Roman, A. R. and Wente, S. R. (2008). Inositol polyphosphates: a new frontier for regulating gene expression. *Chromosoma* **117**, 1–13.

- Allbritton, N. L., Oancea, E., Kuhn, M. A. and Meyer, T. (1994). Source of nuclear calcium signals. *Proc. Natl. Acad. Sci. USA* **91**, 12458-12462.
- Badminton, M. N., Kendall, J. M., Sala-Newby, G. and Campbell, A. K. (1995). Nucleoplasm-targeted aequorin provides evidence for a nuclear calcium barrier. *Exp. Cell Res.* **216**, 236-243.
- Badminton, M. N., Campbell, A. K. and Rembold, C. M. (1996). Differential regulation of nuclear and cytosolic Ca^{2+} in HeLa cells. *J. Biol. Chem.* **271**, 31210-31214.
- Bare, D. J., Kettlun, C. S., Liang, M., Bers, D. M. and Mignery, G. A. (2005). Cardiac type 2 inositol 1,4,5-trisphosphate receptor: interaction and modulation by calcium/calmodulin-dependent protein kinase II. *J. Biol. Chem.* **280**, 15912-15920.
- Bennett, D. L., Cheek, T. R., Berridge, M. J., De Smedt, H., Parys, J. B., Missiaen, L. and Bootman, M. D. (1996). Expression and function of ryanodine receptors in nonexcitable cells. *J. Biol. Chem.* **271**, 6356-6362.
- Berridge, M. J., Lipp, P. and Bootman, M. D. (2000). The versatility and universality of calcium signalling. *Nat. Rev. Mol. Cell Biol.* **1**, 11-21.
- Bezin, S., Charpentier, G., Lee, H. C., Baux, G., Fossier, P. and Cancela, J. M. (2008a). Regulation of nuclear Ca^{2+} signalling by translocation of the Ca^{2+} messenger synthesizing enzyme ADP-ribosyl cyclase during neuronal depolarization. *J. Biol. Chem.* **283**, 27859-27870.
- Bezin, S., Fossier, P. and Cancela, J. M. (2008b). Nucleoplasmic reticulum is not essential in nuclear calcium signalling mediated by cyclic ADPribose in primary neurons. *Pflugers Arch.* **456**, 581-586.
- Birch, B. D., Eng, D. L. and Kocsis, J. D. (1992). Intracellular Ca^{2+} transients during neurite regeneration of an adult mammalian neuron. *Proc. Natl. Acad. Sci. USA* **89**, 7978-7982.
- Bkaily, G., Pothier, P., D'Orleans-Juste, P., Simaan, M., Jacques, D., Jaalouk, D., Belzile, F., Hassan, G., Boutin, C., Haddad, G. et al. (1997). The use of confocal microscopy in the investigation of cell structure and function in the heart, vascular endothelium and smooth muscle cells. *Mol. Cell Biochem.* **172**, 171-194.
- Bkaily, G., Nader, M., Avedanian, L., Choufani, S., Jacques, D., D'Orleans-Juste, P., Gobeil, F., Chemtob, S. and Al-Khoury, J. (2006). G-protein-coupled receptors, channels, and $\text{Na}^+\text{-H}^+$ exchanger in nuclear membranes of heart, hepatic, vascular endothelial, and smooth muscle cells. *Can. J. Physiol. Pharmacol.* **84**, 431-441.
- Bootman, M. D., Thomas, D., Tovey, S. C., Berridge, M. J. and Lipp, P. (2000). Nuclear calcium signalling. *Cell Mol. Life Sci.* **57**, 371-378.
- Bootman, M. D., Collins, T. J., Peppiatt, C. M., Prothero, L. S., MacKenzie, L., De Smet, P., Travers, M., Tovey, S. C., Seo, J. T., Berridge, M. J. et al. (2001). Calcium signalling—an overview. *Semin. Cell Dev. Biol.* **12**, 3-10.
- Bootman, M. D., Berridge, M. J. and Roderick, H. L. (2002). Calcium signalling: more messengers, more channels, more complexity. *Curr. Biol.* **12**, R563-R565.
- Bootman, M. D., Harzheim, D., Smyrniak, I., Conway, S. J. and Roderick, H. L. (2007). Temporal changes in atrial EC-coupling during prolonged stimulation with endothelin-1. *Cell Calcium* **42**, 489-501.
- Brini, M., Murgia, M., Pasti, L., Picard, D., Pozzan, T. and Rizzuto, R. (1993). Nuclear Ca^{2+} concentration measured with specifically targeted recombinant aequorin. *EMBO J.* **12**, 4813-4819.
- Bustamante, J. O., Michelette, E. R., Geibel, J. P., Dean, D. A., Hanover, J. A. and McDonnell, T. J. (2000). Calcium, ATP and nuclear pore channel gating. *Pflugers Arch.* **439**, 433-444.
- Chamero, P., Villalobos, C., Alonso, M. T. and Garcia-Sancho, J. (2002). Dampening of cytosolic Ca^{2+} oscillations on propagation to nucleus. *J. Biol. Chem.* **277**, 50226-50229.
- Chawla, S., Hardingham, G. E., Quinn, D. R. and Bading, H. (1998). CBP: a signal-regulated transcriptional coactivator controlled by nuclear calcium and CaM kinase IV. *Science* **281**, 1505-1509.
- Cheng, H. Y., Pitcher, G. M., Laviolette, S. R., Whishaw, I. Q., Tong, K. I., Kockeritz, L. K., Wada, T., Joza, N. A., Crackower, M., Goncalves, J. et al. (2002). DREAM is a critical transcriptional repressor for pain modulation. *Cell* **108**, 31-43.
- Cockcroft, S. (2006). The latest phospholipase C, PLCeta, is implicated in neuronal function. *Trends Biochem. Sci.* **31**, 4-7.
- Coffey, R. J., Hawkey, C. J., Damstrup, L., Graves-Deal, R., Daniel, V. C., Dempsey, P. J., Chinery, R., Kirkland, S. C., DuBois, R. N., Jetton, T. L. et al. (1997). Epidermal growth factor receptor activation induces nuclear targeting of cyclooxygenase-2, basolateral release of prostaglandins, and mitogenesis in polarizing colon cancer cells. *Proc. Natl. Acad. Sci. USA* **94**, 657-662.
- Danker, T., Mazzanti, M., Tonini, R., Rakowska, A. and Oberleithner, H. (1997). Using atomic force microscopy to investigate patch-clamped nuclear membrane. *Cell Biol. Int.* **21**, 747-757.
- Danker, T., Shahin, V., Schlune, A., Schafer, C. and Oberleithner, H. (2001). Electrophoretic plugging of nuclear pores by using the nuclear hourglass technique. *J. Membr. Biol.* **184**, 91-99.
- De Santa, F., Albini, S., Mezzaroma, E., Baron, L., Felsani, A. and Caruso, M. (2007). pRb-dependent cyclin D3 protein stabilization is required for myogenic differentiation. *Mol. Cell Biol.* **27**, 7248-7265.
- Deisseroth, K., Heist, E. K. and Tsien, R. W. (1998). Translocation of calmodulin to the nucleus supports CREB phosphorylation in hippocampal neurons. *Nature* **392**, 198-202.
- Divecha, N., Banfic, H. and Irvine, R. F. (1991). The polyphosphoinositide cycle exists in the nuclei of Swiss 3T3 cells under the control of a receptor (for IGF-I) in the plasma membrane, and stimulation of the cycle increases nuclear diacylglycerol and apparently induces translocation of protein kinase C to the nucleus. *EMBO J.* **10**, 3207-3214.
- Dolmetsch, R. E., Xu, K. and Lewis, R. S. (1998). Calcium oscillations increase the efficiency and specificity of gene expression. *Nature* **392**, 933-936.
- Dolmetsch, R. E., Pajvani, U., Fife, K., Spotts, J. M. and Greenberg, M. E. (2001). Signaling to the nucleus by an L-type calcium channel-calmodulin complex through the MAP kinase pathway. *Science* **294**, 333-339.
- Downes, C. P., Gray, A. and Lucocq, J. M. (2005). Probing phosphoinositide functions in signaling and membrane trafficking. *Trends Cell Biol.* **15**, 259-268.
- Echevarria, W., Leite, M. F., Guerra, M. T., Zipfel, W. R. and Nathanson, M. H. (2003). Regulation of calcium signals in the nucleus by a nucleoplasmic reticulum. *Nat. Cell Biol.* **5**, 440-446.
- Eder, A. and Bading, H. (2007). Calcium signals can freely cross the nuclear envelope in hippocampal neurons: somatic calcium increases generate nuclear calcium transients. *BMC Neurosci.* **8**, 57.
- Erickson, E. S., Moore, O. L., Moore, D., Krogmeier, J. R. and Dunn, R. C. (2006). The role of nuclear envelope calcium in modifying nuclear pore complex structure. *Can. J. Physiol. Pharmacol.* **84**, 309-318.
- Faenza, I., Ramazzotti, G., Bavelloni, A., Fiume, R., Gaboardi, G. C., Folio, M. Y., Gilmour, R. S., Martelli, A. M., Ravid, K. and Cocco, L. (2007). Inositide-dependent phospholipase C signaling mimics insulin in skeletal muscle differentiation by affecting specific regions of the cyclin D3 promoter. *Endocrinology* **148**, 1108-1117.
- Faenza, I., Bregoli, L., Ramazzotti, G., Gaboardi, G., Folio, M. Y., Mongiorgi, S., Billi, A. M., Manzoli, L., Martelli, A. M. and Cocco, L. (2008). Nuclear phospholipase C beta1 and cellular differentiation. *Front. Biosci.* **13**, 2452-2463.
- Fedoryak, O. D., Searls, Y., Smirnova, I. V., Burns, D. M. and Stehno-Bittel, L. (2004). Spontaneous Ca^{2+} oscillations in subcellular compartments of vascular smooth muscle cells rely on different Ca^{2+} pools. *Cell Res.* **14**, 379-388.
- Fox, J. L., Burgstahler, A. D. and Nathanson, M. H. (1997). Mechanism of long-range Ca^{2+} signalling in the nucleus of isolated rat hepatocytes. *Biochem. J.* **326**, 491-495.
- Fricker, M., Hollinshead, M., White, N. and Vaux, D. (1997). Interphase nuclei of many mammalian cell types contain deep, dynamic, tubular membrane-bound invaginations of the nuclear envelope. *J. Cell Biol.* **136**, 531-544.
- Garcia, K. D., Shah, T. and Garcia, J. (2004). Immunolocalization of type 2 inositol 1,4,5-trisphosphate receptors in cardiac myocytes from newborn mice. *Am. J. Physiol. Cell Physiol.* **287**, C1048-C1057.
- Garner, M. H. (2002). Na,K-ATPase in the nuclear envelope regulates Na^+ : K^+ gradients in hepatocyte nuclei. *J. Membr. Biol.* **187**, 97-115.
- Gerasimenko, J., Maruyama, Y., Tepikin, A., Petersen, O. H. and Gerasimenko, O. (2003). Calcium signalling in and around the nuclear envelope. *Biochem. Soc. Trans.* **31**, 76-78.
- Gerasimenko, O. and Gerasimenko, J. (2004). New aspects of nuclear calcium signalling. *J. Cell Sci.* **117**, 3087-3094.
- Gerasimenko, O. V., Gerasimenko, J. V., Tepikin, A. V. and Petersen, O. H. (1995). ATP-dependent accumulation and inositol trisphosphate- or cyclic ADP-ribose-mediated release of Ca^{2+} from the nuclear envelope. *Cell* **80**, 439-444.
- Gomes, D. A., Leite, M. F., Bennett, A. M. and Nathanson, M. H. (2006). Calcium signaling in the nucleus. *Can. J. Physiol. Pharmacol.* **84**, 325-332.
- Gomes, D. A., Rodrigues, M. A., Leite, M. F., Gomez, M. V., Varnai, P., Balla, T., Bennett, A. M. and Nathanson, M. H. (2008). c-Met must translocate to the nucleus to initiate calcium signals. *J. Biol. Chem.* **283**, 4344-4351.
- Greber, U. F. and Gerace, L. (1992). Nuclear protein import is inhibited by an antibody to a luminal epitope of a nuclear pore complex glycoprotein. *J. Cell Biol.* **116**, 15-30.
- Greber, U. F. and Gerace, L. (1995). Depletion of calcium from the lumen of endoplasmic reticulum reversibly inhibits passive diffusion and signal-mediated transport into the nucleus. *J. Cell Biol.* **128**, 5-14.
- Greber, U. F., Senior, A. and Gerace, L. (1990). A major glycoprotein of the nuclear pore complex is a membrane-spanning polypeptide with a large luminal domain and a small cytoplasmic tail. *EMBO J.* **9**, 1495-1502.
- Guatimosim, S., Amaya, M. J., Guerra, M. T., Aguiar, C. J., Goes, A. M., Gomez-Viquez, N. L., Rodrigues, M. A., Gomes, D. A., Martins-Cruz, J., Lederer, W. J. et al. (2008). Nuclear Ca^{2+} regulates cardiomyocyte function. *Cell Calcium* **44**, 230-242.
- Hallhuber, M., Burkard, N., Wu, R., Buch, M. H., Engelhardt, S., Hein, L., Neyses, L., Schuh, K. and Ritter, O. (2006). Inhibition of nuclear import of calcineurin prevents myocardial hypertrophy. *Circ. Res.* **99**, 626-635.
- Hardingham, G. E., Chawla, S., Johnson, C. M. and Bading, H. (1997). Distinct functions of nuclear and cytoplasmic calcium in the control of gene expression. *Nature* **385**, 260-265.
- Hennager, D. J., Welsh, M. J. and DeLisle, S. (1995). Changes in either cytosolic or nucleoplasmic inositol 1,4,5-trisphosphate levels can control nuclear Ca^{2+} concentration. *J. Biol. Chem.* **270**, 4959-4962.
- Higazi, D. R., Fearnley, C. J., Drawnel, F. M., Talasila, A., Corps, E. M., Ritter, O., McDonald, F., Mikoshiba, K., Bootman, M. D. and Roderick, H. L. (2009). Endothelin-1-stimulated InsP_3 -induced Ca^{2+} release is a nexus for hypertrophic signaling in cardiac myocytes. *Mol. Cell* **33**, 472-482.
- Hsu, A. L., Lu, P. J. and Chen, C. S. (1998). Regulation of nuclear calcium uptake by inositol phosphates and external calcium. *Biochem. Biophys. Res. Commun.* **243**, 653-656.
- Huh, Y. H., Kim, K. D. and Yoo, S. H. (2007). Comparison of and chromogranin effect on inositol 1,4,5-trisphosphate sensitivity of cytoplasmic and nucleoplasmic inositol 1,4,5-trisphosphate receptor/ Ca^{2+} channels. *Biochemistry* **46**, 14032-14043.
- Humbert, J. P., Matter, N., Artault, J. C., Koppler, P. and Malviya, A. N. (1996). Inositol 1,4,5-trisphosphate receptor is located to the inner nuclear membrane vindicating regulation of nuclear calcium signaling by inositol 1,4,5-trisphosphate. Discrete distribution of inositol phosphate receptors to inner and outer nuclear membranes. *J. Biol. Chem.* **271**, 478-485.

- Irvine, R. F. (2006). Nuclear inositolide signalling: expansion, structures and clarification. *Biochim. Biophys. Acta* **1761**, 505-508.
- Ito, M., Shikano, T., Kuroda, K. and Miyazaki, S. (2008). Relationship between nuclear sequestration of PLC ζ and termination of PLC ζ -induced Ca²⁺ oscillations in mouse eggs. *Cell Calcium* **44**, 400-410.
- Jong, Y. J., Kumar, V., Kingston, A. E., Romano, C. and O'Malley, K. L. (2005). Functional metabotropic glutamate receptors on nuclei from brain and primary cultured striatal neurons: role of transporters in delivering ligand. *J. Biol. Chem.* **280**, 30469-30480.
- Kim, C. G., Park, D. and Rhee, S. G. (1996). The role of carboxyl-terminal basic amino acids in Gq α -dependent activation, particulate association, and nuclear localization of phospholipase C- β 1. *J. Biol. Chem.* **271**, 21187-21192.
- Kockskamper, J., Seidlmayer, L., Walther, S., Hellenkamp, K., Maier, L. S. and Pieske, B. (2008a). Endothelin-1 enhances nuclear Ca²⁺ transients in atrial myocytes through Ins(1,4,5)P₃-dependent Ca²⁺ release from perinuclear Ca²⁺ stores. *J. Cell Sci.* **121**, 186-195.
- Kockskamper, J., Zima, A. V., Roderick, H. L., Pieske, B., Blatter, L. A. and Bootman, M. D. (2008b). Emerging roles of inositol 1,4,5-trisphosphate signaling in cardiac myocytes. *J. Mol. Cell Cardiol.* **45**, 128-147.
- Kornhauser, J. M., Cowan, C. W., Shaywitz, A. J., Dolmetsch, R. E., Griffith, E. C., Hu, L. S., Haddad, C., Xia, Z. and Greenberg, M. E. (2002). CREB transcriptional activity in neurons is regulated by multiple, calcium-specific phosphorylation events. *Neuron* **34**, 221-233.
- Kramer, A., Ludwig, Y., Shahin, V. and Oberleithner, H. (2007). A pathway separate from the central channel through the nuclear pore complex for inorganic ions and small macromolecules. *J. Biol. Chem.* **282**, 31437-31443.
- Kumar, V., Jong, Y. J. and O'Malley, K. L. (2008). Activated nuclear metabotropic glutamate receptor mGlu5 couples to nuclear Gq/11 proteins to generate inositol 1,4,5-trisphosphate-mediated nuclear Ca²⁺ release. *J. Biol. Chem.* **283**, 14072-14083.
- Laflamme, K., Domingue, O., Guillemette, B. I. and Guillemette, G. (2002). Immunohistochemical localization of type 2 inositol 1,4,5-trisphosphate receptor to the nucleus of different mammalian cells. *J. Cell Biochem.* **85**, 219-228.
- Larman, M. G., Saunders, C. M., Carroll, J., Lai, F. A. and Swann, K. (2004). Cell cycle-dependent Ca²⁺ oscillations in mouse embryos are regulated by nuclear targeting of PLC ζ . *J. Cell Sci.* **117**, 2513-2521.
- Ledeer, R. and Wu, G. (2007). GM1 in the nuclear envelope regulates nuclear calcium through association with a nuclear sodium-calcium exchanger. *J. Neurochem.* **103 Suppl.** **1**, 126-134.
- Ledeer, R. W. and Wu, G. (2008). Nuclear sphingolipids: metabolism and signaling. *J. Lipid Res.* **49**, 1176-1186.
- Ledo, F., Kremer, L., Mellstrom, B. and Naranjo, J. R. (2002). Ca²⁺-dependent block of CREB-CBP transcription by repressor DREAM. *EMBO J.* **21**, 4583-4592.
- Lee, M. A., Dunn, R. C., Clapham, D. E. and Stehno-Bittel, L. (1998). Calcium regulation of nuclear pore permeability. *Cell Calcium* **23**, 91-101.
- Lee, R. K., Lui, P. P., Ngan, E. K., Lui, J. C., Suen, Y. K., Chan, F. and Kong, S. K. (2006). The nuclear tubular invaginations are dynamic structures inside the nucleus of HeLa cells. *Can. J. Physiol. Pharmacol.* **84**, 477-486.
- Leite, M. F., Thrower, E. C., Echevarria, W., Koulen, P., Hirata, K., Bennett, A. M., Ehrlich, B. E. and Nathanson, M. H. (2003). Nuclear and cytosolic calcium are regulated independently. *Proc. Natl. Acad. Sci. USA* **100**, 2975-2980.
- Limback-Stokin, K., Korzus, E., Nagaoka-Yasuda, R. and Mayford, M. (2004). Nuclear calcium/calmodulin regulates memory consolidation. *J. Neurosci.* **24**, 10858-10867.
- Lipp, P., Thomas, D., Berridge, M. J. and Bootman, M. D. (1997). Nuclear calcium signalling by individual cytoplasmic calcium puffs. *EMBO J.* **16**, 7166-7173.
- Lipp, P., Laine, M., Tovey, S. C., Burrell, K. M., Berridge, M. J., Li, W. and Bootman, M. D. (2000). Functional InsP₃ receptors that may modulate excitation-contraction coupling in the heart. *Curr. Biol.* **10**, 939-942.
- Liu, J., He, Z. Y., Xu, S. M., Liu, F. Y. and Wang, P. Y. (2001). Inositol 1,4,5-trisphosphate receptors (IP₃Rs) in myocardial nuclei involved in pressure overload-induced hypertrophy of rat heart. *Sheng Li Xue Bao* **53**, 281-285.
- Lui, P. P., Kong, S. K., Fung, K. P. and Lee, C. Y. (1998). The rise of nuclear and cytosolic Ca²⁺ can be uncoupled in HeLa cells. *Pflügers Arch.* **436**, 371-376.
- Lui, P. P., Chan, F. L., Suen, Y. K., Kwok, T. T. and Kong, S. K. (2003). The nucleus of HeLa cells contains tubular structures for Ca²⁺ signaling with the involvement of mitochondria. *Biochem. Biophys. Res. Commun.* **308**, 826-833.
- Luo, D., Yang, D., Lan, X., Li, K., Li, X., Chen, J., Zhang, Y., Xiao, R. P., Han, Q. and Cheng, H. (2008). Nuclear Ca²⁺ sparks and waves mediated by inositol 1,4,5-trisphosphate receptors in neonatal rat cardiomyocytes. *Cell Calcium* **43**, 165-174.
- Luo, D. L., Gao, J., Lan, X. M., Wang, G., Wei, S., Xiao, R. P. and Han, Q. D. (2006). Role of inositol 1,4,5-trisphosphate receptors in α 1-adrenergic receptor-induced cardiomyocyte hypertrophy. *Acta Pharmacol. Sin.* **27**, 895-900.
- Malviya, A. N. (1994). The nuclear inositol 1,4,5-trisphosphate and inositol 1,3,4,5-tetrakisphosphate receptors. *Cell Calcium* **16**, 301-313.
- Malviya, A. N. and Klein, C. (2006). Mechanism regulating nuclear calcium signaling. *Can. J. Physiol. Pharmacol.* **84**, 403-422.
- Marius, P., Guerra, M. T., Nathanson, M. H., Ehrlich, B. E. and Leite, M. F. (2006). Calcium release from ryanodine receptors in the nucleoplasmic reticulum. *Cell Calcium* **39**, 65-73.
- Mellstrom, B., Savignac, M., Gomez-Villafuertes, R. and Naranjo, J. R. (2008). Ca²⁺-operated transcriptional networks: molecular mechanisms and *in vivo* models. *Physiol. Rev.* **88**, 421-449.
- Mermelstein, P. G., Deisseroth, K., Dasgupta, N., Isaksen, A. L. and Tsien, R. W. (2001). Calmodulin priming: nuclear translocation of a calmodulin complex and the memory of prior neuronal activity. *Proc. Natl. Acad. Sci. USA* **98**, 15342-15347.
- Molkentin, J. D. (2006). Dichotomy of Ca²⁺ in the heart: contraction versus intracellular signaling. *J. Clin. Invest.* **116**, 623-626.
- O'Brien, E. M., Gomes, D. A., Sehgal, S. and Nathanson, M. H. (2007). Hormonal regulation of nuclear permeability. *J. Biol. Chem.* **282**, 4210-4217.
- O'Malley, D. M., Burbach, B. J. and Adams, P. R. (1999). Fluorescent calcium indicators: subcellular behavior and use in confocal imaging. *Methods Mol. Biol.* **122**, 261-303.
- Okada, M., Ishimoto, T., Naito, Y., Hirata, H. and Yagisawa, H. (2005). Phospholipase C δ 1 associates with importin β 1 and translocates into the nucleus in a Ca²⁺-dependent manner. *FEBS Lett.* **579**, 4949-4954.
- Perez-Terzic, C., Stehno-Bittel, L. and Clapham, D. E. (1997). Nucleoplasmic and cytoplasmic differences in the fluorescence properties of the calcium indicator Fluo-3. *Cell Calcium* **21**, 275-282.
- Perez-Terzic, C., Gacy, A. M., Bortolon, R., Dzeja, P. P., Puceat, M., Jaconi, M., Prendergast, F. G. and Terzic, A. (1999). Structural plasticity of the cardiac nuclear pore complex in response to regulators of nuclear import. *Circ. Res.* **84**, 1292-1301.
- Perez, P. J., Ramos-Franco, J., Fill, M. and Mignery, G. A. (1997). Identification and functional reconstitution of the type 2 inositol 1,4,5-trisphosphate receptor from ventricular cardiac myocytes. *J. Biol. Chem.* **272**, 23961-23969.
- Pust, T., Wu, J. J., Zimmerman, T. L., Zhang, L., Ehrlich, B. E., Berchtold, M. W., Hoek, J. B., Karpen, S. J., Nathanson, M. H. and Bennett, A. M. (2002). Epidermal growth factor-mediated activation of the ETS domain transcription factor Elk-1 requires nuclear calcium. *J. Biol. Chem.* **277**, 27517-27527.
- Quesada, I. and Verdugo, P. (2005). InsP₃ signaling induces pulse-modulated Ca²⁺ signals in the nucleus of airway epithelial ciliated cells. *Biophys. J.* **88**, 3946-3953.
- Ramazzotti, G., Faenza, I., Gaboardi, G. C., Piazzi, M., Bavelloni, A., Fiume, R., Manzoli, L., Martelli, A. M. and Cocco, L. (2008). Catalytic activity of nuclear PLC- β 1 is required for its signalling function during C2C12 differentiation. *Cell. Signal.* **22**, 2013-2021.
- Roderick, H. L. and Bootman, M. D. (2003). Bi-directional signalling from the InsP₃ receptor: regulation by calcium and accessory factors. *Biochem. Soc. Trans.* **31**, 950-953.
- Roderick, H. L., Higazi, D. R., Smyrniak, I., Fearnley, C., Harzheim, D. and Bootman, M. D. (2007). Calcium in the heart: when it's good, it's very very good, but when it's bad, it's horrid. *Biochem. Soc. Trans.* **35**, 957-961.
- Rodrigues, M. A., Gomes, D. A., Leite, M. F., Grant, W., Zhang, L., Lam, W., Cheng, Y. C., Bennett, A. M. and Nathanson, M. H. (2007). Nucleoplasmic calcium is required for cell proliferation. *J. Biol. Chem.* **282**, 17061-17068.
- Rodrigues, M. A., Gomes, D. A., Andrade, V. A., Leite, M. F. and Nathanson, M. H. (2008). Insulin induces calcium signals in the nucleus of rat hepatocytes. *Hepatology* **48**, 1621-1631.
- Santella, L. and Carafoli, E. (1997). Calcium signaling in the cell nucleus. *FASEB J.* **11**, 1091-1109.
- Santella, L., Ercolano, E., Lim, D., Nusco, G. A. and Moccia, F. (2003). Activated M-phase-promoting factor (MPF) is exported from the nucleus of starfish oocytes to increase the sensitivity of the Ins(1,4,5)P₃ receptors. *Biochem. Soc. Trans.* **31**, 79-82.
- Savard, M., Barbaz, D., Belanger, S., Muller-Esterl, W., Bkaily, G., D'Orleans-Juste, P., Cote, J., Bovenzi, V. and Gobeil, F., Jr (2008). Expression of endogenous nuclear bradykinin B2 receptors mediating signaling in immediate early gene activation. *J. Cell Physiol.* **216**, 234-244.
- Schermelleh, L., Carlton, P. M., Haase, S., Shao, L., Winoto, L., Kner, P., Burke, B., Cardoso, M. C., Agard, D. A., Gustafsson, M. G. et al. (2008). Subdiffraction multicolor imaging of the nuclear periphery with 3D structured illumination microscopy. *Science* **320**, 1332-1336.
- Schievella, A. R., Regier, M. K., Smith, W. L. and Lin, L. L. (1995). Calcium-mediated translocation of cytosolic phospholipase A₂ to the nuclear envelope and endoplasmic reticulum. *J. Biol. Chem.* **270**, 30749-30754.
- Shahin, V., Danker, T., Enss, K., Ossig, R. and Oberleithner, H. (2001). Evidence for Ca²⁺- and ATP-sensitive peripheral channels in nuclear pore complexes. *FASEB J.* **15**, 1895-1901.
- Shaw, J. P., Utz, P. J., Durand, D. B., Toole, J. J., Emmel, E. A. and Crabtree, G. R. (1988). Identification of a putative regulator of early T cell activation genes. *Science* **241**, 202-205.
- Shaywitz, A. J. and Greenberg, M. E. (1999). CREB: a stimulus-induced transcription factor activated by a diverse array of extracellular signals. *Annu. Rev. Biochem.* **68**, 821-861.
- Shibasaki, F., Price, E. R., Milan, D. and McKeon, F. (1996). Role of kinases and the phosphatase calcineurin in the nuclear shuttling of transcription factor NF-AT4. *Nature* **382**, 370-373.
- Shirakawa, H. and Miyazaki, S. (1996). Spatiotemporal analysis of calcium dynamics in the nucleus of hamster oocytes. *J. Physiol.* **494**, 29-40.
- Soares, S., Thompson, M., White, T., Isbell, A., Yamasaki, M., Prakash, Y., Lund, F. E., Galione, A. and Chini, E. N. (2007). NAADP as a second messenger: neither CD38 nor base-exchange reaction are necessary for *in vivo* generation of NAADP in myometrial cells. *Am. J. Physiol. Cell Physiol.* **292**, C227-C239.
- Stallings, J. D., Zeng, Y. X., Narvaez, F. and Rebecchi, M. J. (2008). Phospholipase C- δ 1 expression is linked to proliferation, DNA synthesis, and cyclin E levels. *J. Biol. Chem.* **283**, 13992-14001.

- Stehno-Bittel, L., Luckhoff, A. and Clapham, D. E. (1995). Calcium release from the nucleus by InsP3 receptor channels. *Neuron* **14**, 163-167.
- Stoffler, D., Goldie, K. N., Feja, B. and Aebi, U. (1999). Calcium-mediated structural changes of native nuclear pore complexes monitored by time-lapse atomic force microscopy. *J. Mol. Biol.* **287**, 741-752.
- Stoffler, D., Schwarz-Herion, K., Aebi, U. and Fahrenkrog, B. (2006). Getting across the nuclear pore complex: new insights into nucleocytoplasmic transport. *Can. J. Physiol. Pharmacol.* **84**, 499-507.
- Sugden, P. H. and Clerk, A. (2005). Endothelin signalling in the cardiac myocyte and its pathophysiological relevance. *Curr. Vasc. Pharmacol.* **3**, 343-351.
- Suh, P. G., Park, J. I., Manzoli, L., Cocco, L., Peak, J. C., Katan, M., Fukami, K., Kataoka, T., Yun, S. and Ryu, S. H. (2008). Multiple roles of phosphoinositide-specific phospholipase C isozymes. *BMB Rep.* **41**, 415-434.
- Thomas, D., Lipp, P., Tovey, S. C., Berridge, M. J., Li, W., Tsien, R. Y. and Bootman, M. D. (2000a). Microscopic properties of elementary Ca^{2+} release sites in non-excitable cells. *Curr. Biol.* **10**, 8-15.
- Thomas, D., Tovey, S. C., Collins, T. J., Bootman, M. D., Berridge, M. J. and Lipp, P. (2000b). A comparison of fluorescent Ca^{2+} indicator properties and their use in measuring elementary and global Ca^{2+} signals. *Cell Calcium* **28**, 213-223.
- Thorogate, R. and Torok, K. (2004). Ca^{2+} -dependent and -independent mechanisms of calmodulin nuclear translocation. *J. Cell Sci.* **117**, 5923-5936.
- Thorogate, R. and Torok, K. (2007). Role of Ca^{2+} activation and bilobal structure of calmodulin in nuclear and nucleolar localization. *Biochem. J.* **402**, 71-80.
- Thrower, E. C., Choe, C. U., So, S. H., Jeon, S. H., Ehrlich, B. E. and Yoo, S. H. (2003). A functional interaction between chromogranin B and the inositol 1,4,5-trisphosphate receptor/ Ca^{2+} channel. *J. Biol. Chem.* **278**, 49699-49706.
- Tomida, T., Hirose, K., Takizawa, A., Shibasaki, F. and Iino, M. (2003). NFAT functions as a working memory of Ca^{2+} signals in decoding Ca^{2+} oscillation. *EMBO J.* **22**, 3825-3832.
- Torok, K. (2007). The regulation of nuclear membrane permeability by Ca^{2+} signaling: a tightly regulated pore or a floodgate? *Sci. STKE* 2007, pe24.
- Trubiani, O., Guarnieri, S., Eleuterio, E., Di Giuseppe, F., Orciani, M., Angelucci, S. and Di Primio, R. (2008). Insights into nuclear localization and dynamic association of CD38 in Raji and K562 cells. *J. Cell Biochem.* **103**, 1294-1308.
- Vermassen, E., Van Acker, K., Annaert, W. G., Himpens, B., Callewaert, G., Missiaen, L., De Smedt, H. and Parys, J. B. (2003). Microtubule-dependent redistribution of the type-1 inositol 1,4,5-trisphosphate receptor in A7r5 smooth muscle cells. *J. Cell Sci.* **116**, 1269-1277.
- Visnjic, D. and Banfic, H. (2007). Nuclear phospholipid signaling: phosphatidylinositol-specific phospholipase C and phosphoinositide 3-kinase. *Pflugers Arch.* **455**, 19-30.
- Wang, H. and Clapham, D. E. (1999). Conformational changes of the in situ nuclear pore complex. *Biophys. J.* **77**, 241-247.
- Wei, X., Henke, V. G., Strubing, C., Brown, E. B. and Clapham, D. E. (2003). Real-time imaging of nuclear permeation by EGFP in single intact cells. *Biophys. J.* **84**, 1317-1327.
- Wu, X. and Bers, D. M. (2006). Sarcoplasmic reticulum and nuclear envelope are one highly interconnected Ca^{2+} store throughout cardiac myocyte. *Circ. Res.* **99**, 283-291.
- Wu, X. and Bers, D. M. (2007). Free and bound intracellular calmodulin measurements in cardiac myocytes. *Cell Calcium* **41**, 353-364.
- Wu, X., Zhang, T., Bossuyt, J., Li, X., McKinsey, T. A., Dedman, J. R., Olson, E. N., Chen, J., Brown, J. H. and Bers, D. M. (2006). Local InsP3-dependent perinuclear Ca^{2+} signaling in cardiac myocyte excitation-transcription coupling. *J. Clin. Invest.* **116**, 675-682.
- Xie, X., Wu, G., Lu, Z. H. and Ledeen, R. W. (2002). Potentiation of a sodium-calcium exchanger in the nuclear envelope by nuclear GM1 ganglioside. *J. Neurochem.* **81**, 1185-1195.
- Xie, X., Wu, G. and Ledeen, R. W. (2004). C6 cells express a sodium-calcium exchanger/GM1 complex in the nuclear envelope but have no exchanger in the plasma membrane: comparison to astrocytes. *J. Neurosci. Res.* **76**, 363-375.
- Yagisawa, H. (2006). Nucleocytoplasmic shuttling of phospholipase C- δ 1: a link to Ca^{2+} . *J. Cell Biochem.* **97**, 233-243.
- Ye, K. and Ahn, J. Y. (2008). Nuclear phosphoinositide signaling. *Front. Biosci.* **13**, 540-548.
- Yoo, S. H., Chu, S. Y., Kim, K. D. and Huh, Y. H. (2007). Presence of secretogranin II and high-capacity, low-affinity Ca^{2+} storage role in nucleoplasmic Ca^{2+} store vesicles. *Biochemistry* **46**, 14663-14671.
- Zima, A. V., Bare, D. J., Mignery, G. A. and Blatter, L. A. (2007). IP3-dependent nuclear Ca^{2+} signalling in the mammalian heart. *J. Physiol.* **584**, 601-611.

Commentaries and Cell Science at a Glance

JCS Commentaries highlight and critically discuss recent and exciting findings that will interest those who work in cell biology, molecular biology, genetics and related disciplines, whereas Cell Science at a Glance poster articles are short primers that act as an introduction to an area of cell biology, and include a large poster and accompanying text.

Both of these article types, designed to appeal to specialists and nonspecialists alike, are commissioned from leading figures in the field and are subject to rigorous peer-review and in-house editorial appraisal. Each issue of the journal usually contains at least one of each article type. JCS thus provides readers with more than 50 topical pieces each year, which cover the complete spectrum of cell science. The following are just some of the areas that will be covered in JCS over the coming months:

Cell Science at a Glance

- WASP and SCAR/WAVE proteins – the drivers of actin assembly *Robert Insall*
- Collective cell migration at a glance *Peter Friedl*
- Podosomes and invadopodia at a glance *Stefan Linder*
- Mammalian cytosine methylation at a glance *Timothy Bestor*
- Dynamins at a glance *Jenny Hinshaw*
- mTOR signalling at a glance *David Sabatini*

Commentaries

- The Crumbs complex *Elisabeth Knust*
- How do anti-mitotic drugs kill cancer cells? *Stephen Taylor*
- Mechanisms for transcellular diapedesis *Chris Carman*
- ABL-family kinases *Tony Koleske*
- Tail-anchored protein biogenesis – the beginning for the end? *Stephen High*

Although we discourage the submission of unsolicited Commentaries and Cell Science at a Glance poster articles to the journal, ideas for future articles – in the form of a short proposal and some key references – are welcome and should be sent by email to the Editorial Office (jcs@biologists.com).

Journal of Cell Science, Bidder Building, 140 Cowley Road, Cambridge CB4 0DL, UK
E-mail: jcs@biologists.com Website: <http://jcs.biologists.org>